

**Lucrări științifice (selecție 2013-2023):**

1. A computational study of the steric strain of –N=O groups in saturated nitrosamines, F. Potmischil, M.Hillebrand, *Rev. Roum. Chimie*, **68**, 91 (2023).
2. A New Nitronyl-Nitroxide Ligand for Designing Binuclear Ln III Complexes: Syntheses, Crystal Structures, Magnetic and EPR Studies, M. Raduca, D.O.T. A. Martins, C.A.Spanu, M. Hillebrand, F. Tuna, G.Ionita, A.M.Madalan, C. Lecourt, J.P. Sutter, M.Andruh, *EurJIC* (2022), [doi.org/10.1002/ejic.202200128](https://doi.org/10.1002/ejic.202200128)
3. Synthesis, crystal structure, magnetic, spectroscopic, and theoretical investigations of two new nitronyl-nitroxide complexes, C. A. Spanu, C. Pichon, G. Ionita, T. Mocanu, S. Calancea, M.Raduta, J.P. Sutter, M.Hillebrand, M.Andruh, *J. Coord.Chem.*, **74**, 279 (2021).
4. [Hydroacridines: Part 33. An experimental and DFT study of the 13 C NMR chemical shifts of the nitrosamines derived from the six stereoisomers of tetradecahydroacridine.](#) F. Potmischil, M. Hillebrand, H. Kalchhauser, *Magnetic resonance in chemistry, MRC*, **58**, 3, 2049 (2020).
5. Trinuclear Nickel(II) and Cobalt(II) Complexes Constructed from Mannich–Schiff-Base Ligands: Synthesis, Crystal Structures, and Magnetic Properties, M. I. Mocanu, A. A. Patrascu, M.Hillebrand, S. Shova, F. Lloret, M. Julve and M. Andruh, *Eur. J. Inorg. Chem.*, 4773 (2019).
6. Synthesis, crystal structure, magnetic and catalytic properties of a new dissymmetric binuclear nickel(II) complex, V. A. Neacșu, C. Maxim, M. Hillebrand, M. del Camino González-Arellano, S. Soriano, E. Rentschler, M. Andruh, *Polyhedron*, **150**, 77 (2018).
7. A novel 2-D coordination polymer with mixed azido and alkoxido bridges: Synthesis, structure and magnetic properties. , L.Stoicescu, C. Maxim, M.Rouzières, M. Hillebrand, R. Clérac, M. Andruh, *Polyhedron*, **92**,111 (2015).
8. [Photophysical properties of two derivatives in the phenoxathiin-10,10-dioxide class](#), M. Duca, I. Matei, M. Hillebrand, *Rev Roum. Chim.*, **59**, 683 (2014).
9. Close-lying pKa values of kaempferol determined by second-derivative synchronous fluorescence, I. Matei, C.Tablet, S. Ionescu, M.Hillebrand, *Rev Roum. Chim.*, **59**, 401 (2014).
10.  $\beta$ -Cyclodextrin derivative as chiral carrier in membrane transport of some aromatic amino acids, A. Stancu, M. Hillebrand, C. Tablet, L. Mutihac, *J. Incl. Phenom. Macrocycl Chem*, **78**, 71 (2014).

**11.** [Magnetic and luminescent binuclear double-stranded helicates](#), P. Cucos, F. Tuna, L. Sorace, I. Matei, C. Maxim, S. Shova, R. Gheorghe, A. Caneschi, M. Hillebrand, M. Andruh, *Inorg. Chem.* **53**, 7738 (2014).

**12.** Induced Chirality of Genistein upon Binding to Albumin: Circular Dichroism and TDDFT study, I. Matei, S. Ionescu, M. Hillebrand, *Rev. Roum. Chim.* **58** (2013).

**13.** Phenoxathiinsulphone derivatives–cyclodextrin interactions: induced chirality and TDDFT calculations, R. Sandu, C. Tablet, M. Hillebrand, *J. Incl. Phenom. Macrocycl Chem* **77**, 183 (2013).

**14.** New Insights on Flavonoid-Serum Albumin Interactions from Concerted Spectroscopic Methods and Molecular Modeling, S. Ionescu, I. Matei, C. Tablet, M. Hillebrand; *Current Drug Metabolism* **14** 474, (2013).

**15.** Exploring the capabilities of TDDFT calculations to explain the induced chirality upon a binding process: A simple case, 3-carboxycoumarin, A. Varlan, M. Hillebrand, *J. Mol. Struct.* **1036** 341, (2013).

**16.** Theoretical ECD calculations – a useful tool for estimating the conformational change of a ligand in the binding pocket of proteins, S. Ionescu, I. Matei, C. Tablet, M. Hillebrand, *Phys. Chem. Chem. Phys.* **15** 11604, (2013).