

**Lucrări științifice:**

1. Crystal and molecular structure of tetramethyldithioimidodiphosphinic acid,  $(\text{SPMe}_2)_2\text{NH}$ , and its cobalt(II) complex,  $\text{Co}[(\text{SPMe}_2)_2\text{N}_2]_2$ , containing a tetrahedral  $\text{CoS}_4$  core, împreună cu R. Rösler, I. Haiduc, R. Cea-Olivares and G. Espinosa-Perez, *Inorg. Chem.*, **1995**, 34, 3352-3354, FI ISI / 2007 = 4.123
2. Structural patterns in inorganic and organoantimony derivatives of oxo- and thiodiorganophosphorus ligands, împreună cu I. Haiduc, *Coord. Chem. Rev.*, **1996**, 147, 117-146, FI ISI / 2007 = 8.568
3. The first crystal structures of mixed chalcogen derivatives,  $\text{R}_2\text{Sn}[(\text{OPPh}_2)(\text{SPPh}_2)_2\text{N}]_2$  ( $\text{R} = \text{Me, Ph}$ ), împreună cu R. Rösler, J. E. Drake, J. Yang and I. Haiduc, *J. Chem. Soc., Dalton Trans.*, **1996**, 391-399, FI ISI / 2007 = 3.212
4. Structural chemistry of bismuth compounds. I. Organobismuth derivatives, împreună cu H. J. Breunig and H. Althaus, *Chem. Rev.*, **1999**, 99, 3277-3328, FI ISI / 2007 = 22.757
5. Tetraorganodichalcogenoimidodiphosphorus acids and their Main Group metal derivatives, împreună cu J. E. Drake, *Coord. Chem. Rev.*, **2001**, 223 (1), 117-216, FI ISI / 2007 = 8.568
6. Syntheses and chemistry of hypervalent *cyclo-R*<sub>4</sub>Sb<sub>4</sub>, *cyclo-(RSbE)*<sub>n</sub> [ $\text{R} = 2\text{-}(\text{Me}_2\text{NCH}_2)\text{C}_6\text{H}_4$ , E = O, S] and precursors, împreună cu împreună cu L. M. Opris, A. Silvestru, H. J. Breunig and E. Lork, *Dalton Trans.*, **2004**, 3575 – 3585, FI ISI / 2007 = 3.212
7. Phosphorescent excited state of  $[\text{Au}_2\{(\text{Ph}_2\text{Sb})_2\text{O}\}_3]^{2+}$ : Jahn-Teller distortion at only one gold (I) center, împreună cu V. R. Bojan, E. J. Fernandez, A. Laguna, J. M. Lopez-de-Luzuriaga, M. Monge, M. E. Olmos, *J. Am Chem. Soc.*, **2005**, 127, 11564-11565, FI ISI / 2007 = 7.885
8. Hypercoordinated organotin(IV) halides containing  $2\text{-}(\text{Me}_2\text{NCH}_2)\text{C}_6\text{H}_4$  groups:  $[\text{2-}(\text{Me}_2\text{NCH}_2)\text{C}_6\text{H}_4]_2\text{SnX}_2$  (X = F, Cl, Br, I) and  $[\text{2-}(\text{Me}_2\text{NCH}_2)\text{C}_6\text{H}_4]\text{R}_2\text{SnX}$  ( $\text{R} = \text{Me, Ph}$ ; X = F, Cl, Br, I) – solution behaviour and solid state hydrogen bonding-based supramolecular architecture, împreună cu R. A. Varga, A. Rotar, M. Schuermann, K. Jurkschat, *Eur. J. Inorg. Chem.*, **2006**, 1475-1486, FI ISI / 2007 = 2.597
9. Isomers of a dibismuthane,  $\text{R}_2\text{Bi-BiR}_2$  [ $\text{R} = 2,6\text{-}(\text{Me}_2\text{NCH}_2)_2\text{C}_6\text{H}_3$ ], and unusual reactions with oxygen: formation of  $[\text{R}_2\text{Bi}]_2(\text{O}_2)$  and  $\text{R}'\text{R}''\text{Bi}$  [ $\text{R}' = 2\text{-}(\text{Me}_2\text{NCH}_2)\text{-}6\text{-}\{\text{Me}_2\text{N}(\text{O})\text{CH}_2\}\text{C}_6\text{H}_3$ ;  $\text{R}'' = 2\text{-}(\text{Me}_2\text{NCH}_2)\text{-}6\text{-}\{\text{O}(\text{O})\text{C}\}\text{C}_6\text{H}_3$ ], împreună cu L. Balázs, H. J. Breunig, E Lork, A. Soran, *Inorg. Chem.*, **2006**, 45, 2341-2346, FI ISI / 2007 = 4.123
10. Organobismuth(III) dihalides with T-shaped geometry stabilized by intramolecular N→Bi interactions and related diorganobismuth (III) halides, împreună cu A. P. Soran, H. J. Breunig, G. Balázs and J. C. Green, *Organometallics*, **2007**, 26, 1196-1203, FI ISI / 2007 = 3.833
11. Hypervalent organobismuth(III) carbonate, chalcogenides and halides with the pendant arm ligands  $2\text{-}(\text{Me}_2\text{NCH}_2)\text{C}_6\text{H}_4$  and  $2,6\text{-}(\text{Me}_2\text{NCH}_2)_2\text{C}_6\text{H}_3$ , împreună cu H. J. Breunig, L. Königsmann, E. Lork, N. Philipp, M. Nema, A. Soran, R. A. Varga and R. Wagner, *Dalton Trans.*, **2008**, 1831-1842, FI ISI / 2007 = 3.212
12. Gold-Heterometal Interactions and Bonds, in *Modern Supramolecular Gold Chemistry: Gold-Metal Interactions and Applications*, A. Laguna (Ed.), Wiley-VCH, New York – Weinheim, **2008**, Chapter 4, p. 181-293, ISBN 978-3-527-32029-5