



# COMPOSITIO MATHEMATICA

## An injectivity theorem

Florin Ambro

Compositio Math. **150** (2014), 999–1023.

[doi:10.1112/S0010437X13007768](https://doi.org/10.1112/S0010437X13007768)





# An injectivity theorem

Florin Ambro

## ABSTRACT

We generalize the injectivity theorem of Esnault and Viehweg, and apply it to the structure of log canonical type divisors.

## Introduction

We are interested in the following *lifting problem*: given a Cartier divisor  $L$  on a complex variety  $X$  and a closed subvariety  $Y \subset X$ , when is the restriction map

$$\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L))$$

surjective? The standard method is to consider the short exact sequence

$$0 \rightarrow \mathcal{I}_Y(L) \rightarrow \mathcal{O}_X(L) \rightarrow \mathcal{O}_Y(L) \rightarrow 0,$$

which induces a long exact sequence in cohomology

$$0 \rightarrow \Gamma(X, \mathcal{I}_Y(L)) \rightarrow \Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L)) \rightarrow H^1(X, \mathcal{I}_Y(L)) \xrightarrow{\alpha} H^1(X, \mathcal{O}_X(L)) \cdots$$

The restriction is surjective if and only if  $\alpha$  is injective. In particular, if  $H^1(X, \mathcal{I}_Y(L)) = 0$ .

If  $X$  is a non-singular proper curve, Serre duality answers completely the lifting problem: the restriction map is not surjective if and only if  $L \sim K_X + Y - D$  for some effective divisor  $D$  such that  $D - Y$  is not effective. In particular,  $\deg L \leq \deg(K_X + Y)$ . If  $\deg L > \deg(K_X + Y)$ , then  $H^1(X, \mathcal{I}_Y(L)) = 0$ , and therefore lifting holds.

If  $X$  is a non-singular projective surface, only sufficient criteria for lifting are known (see [Zar35]). If  $H$  is a general hyperplane section induced by a Veronese embedding of sufficiently large degree (depending on  $L$ ), then  $\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(H, \mathcal{O}_H(L))$  is an isomorphism (Enriques–Severi–Zariski). If  $H$  is a hyperplane section of  $X$ , then  $H^i(X, \mathcal{O}_X(K_X + H)) = 0$  ( $i > 0$ ) (Picard–Severi).

These classical results were extended by Serre [Ser55] as follows: if  $X$  is affine and  $\mathcal{F}$  is a quasi-coherent  $\mathcal{O}_X$ -module, then  $H^i(X, \mathcal{F}) = 0$  ( $i > 0$ ). If  $X$  is projective,  $H$  is ample and  $\mathcal{F}$  is a coherent  $\mathcal{O}_X$ -module, then  $H^i(X, \mathcal{F}(mH)) = 0$  ( $i > 0$ ) for  $m$  sufficiently large.

Kodaira [Kod53] extended Picard–Severi’s result as follows: if  $X$  is a projective complex manifold, and  $H$  is an ample divisor, then  $H^i(X, \mathcal{O}_X(K_X + H)) = 0$  ( $i > 0$ ). This vanishing remains true over a field of characteristic zero, but may fail in positive characteristic (Raynaud [Ray78]). Kodaira’s vanishing is central in the classification theory of complex algebraic varieties, but one has to weaken the positivity of  $H$  to apply it successfully: it still holds if  $H$  is

---

Received 27 September 2012, accepted in final form 26 September 2013, published online 12 May 2014.

2010 Mathematics Subject Classification 14F17 (primary), 14E30 (secondary).

Keywords: logarithmic differential forms, log varieties, totally canonical locus, non-log canonical locus.

This work was supported by a grant of the Romanian National Authority for Scientific Research, CNCS – UEFISCDI, project number PN-II-RU-TE-2011-3-0097.

This journal is © Foundation Compositio Mathematica 2014.

only semiample and big (Mumford [Mum67], Ramanujam [Ram72]), or if  $K_X + H$  is replaced by  $\lceil K_X + H \rceil$  for a  $\mathbb{Q}$ -divisor  $H$  which is nef and big, whose fractional part is supported by a normal crossings divisor (Ramanujam [Ram74], Miyaoka [Miy79], Kawamata [Kaw82], Viehweg [Vie82]). Recall that the round up of a real number  $x$  is  $\lceil x \rceil = \min\{n \in \mathbb{Z}; x \leq n\}$ , and the round up of a  $\mathbb{Q}$ -divisor  $D = \sum_E d_E E$  is  $\lceil D \rceil = \sum_E \lceil d_E \rceil E$ .

The first lifting criterion in the absence of bigness is due to Tankeev [Tan71]: if  $X$  is proper non-singular and  $Y \subset X$  is the general member of a free linear system, then the restriction

$$\Gamma(X, \mathcal{O}_X(K_X + 2Y)) \rightarrow \Gamma(Y, \mathcal{O}_Y(K_X + 2Y))$$

is surjective. Kollár [Kol86] extended it to the following injectivity theorem: if  $H$  is a semiample divisor and  $D \in |m_0 H|$  for some  $m_0 \geq 1$ , then the homomorphism

$$H^q(X, \mathcal{O}_X(K_X + mH)) \rightarrow H^q(X, \mathcal{O}_X(K_X + mH + D))$$

is injective for all  $m \geq 1, q \geq 0$ . Esnault and Viehweg [EV86, EV92] removed completely the positivity assumption, to obtain the following injectivity result: let  $L$  be a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{Q}} K_X + \sum_i b_i E_i$ , where  $\sum_i E_i$  is a normal crossings divisor and  $0 \leq b_i \leq 1$  are rational numbers. If  $D$  is an effective divisor supported by  $\sum_{0 < b_i < 1} E_i$ , then the homomorphism

$$H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(X, \mathcal{O}_X(L + D))$$

is injective, for all  $q$ . The original result [EV92, Theorem 5.1] was stated in terms of roots of sections of powers of line bundles, and restated in this logarithmic form in [Amb06, Corollary 3.2]. It was used in [Amb03, Amb06] to derive basic properties of log varieties and quasi-log varieties.

The main result of this paper (Theorem 2.3) is that Esnault–Viehweg’s injectivity remains true even if some components  $E_i$  of  $D$  have  $b_i = 1$ . In fact, it reduces to the special case when all  $b_i = 1$ , which has the following geometric interpretation.

**THEOREM 0.1.** *Let  $X$  be a proper non-singular variety, defined over an algebraically closed field of characteristic zero. Let  $\Sigma$  be a normal crossings divisor on  $X$ , let  $U = X \setminus \Sigma$ . Then the restriction homomorphism*

$$H^q(X, \mathcal{O}_X(K_X + \Sigma)) \rightarrow H^q(U, \mathcal{O}_U(K_U))$$

*is injective, for all  $q$ .*

Combined with Serre vanishing on affine varieties, it gives the following corollary.

**COROLLARY 0.2.** *Let  $X$  be a proper non-singular variety, defined over an algebraically closed field of characteristic zero. Let  $\Sigma$  be a normal crossings divisor on  $X$  such that  $X \setminus \Sigma$  is contained in an affine open subset of  $X$ . Then*

$$H^q(X, \mathcal{O}_X(K_X + \Sigma)) = 0$$

*for  $q > 0$ .*

If  $X \setminus \Sigma$  itself is affine, this vanishing is due to Esnault and Viehweg [EV92, p. 5]. It implies the Kodaira vanishing theorem.

We outline the structure of this paper. After some preliminaries in §1, we prove the main injectivity result in §2. The proof is similar to that of Esnault–Viehweg, except that we do not use duality. It is an immediate consequence of the Atiyah–Hodge lemma and Deligne’s degeneration of

the logarithmic Hodge to de Rham spectral sequence. In §3, we obtain some vanishing theorems for sheaves of logarithmic forms of intermediate degree. The results are the same as in [EV92], except that the complement of the boundary is only contained in an affine open subset, instead of being itself affine. They suggest that injectivity may extend to forms of intermediate degree (Question 7.1). In §4, we introduce the *locus of totally canonical singularities* and the *non-log canonical locus* of a log variety. The latter has the same support as the subscheme structure for the non-log canonical locus introduced in [Amb03], but the scheme structures usually differ (see Remark 4.4). In §5, we partially extend the injectivity theorem to the category of log varieties. The open subset to which we restrict is the locus of totally canonical singularities of some log structure. We can only prove the injectivity for the first cohomology group. The idea is to descend injectivity from a log resolution, and to make this work for higher cohomology groups one needs vanishing theorems or at least the degeneration of the Leray spectral sequence for a certain resolution. We do not pursue this here. In §6, we establish the *lifting property* of  $\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L))$  for a Cartier divisor  $L \sim_{\mathbb{R}} K_X + B$ , with  $Y$  the non-log canonical locus of  $X$  (Theorem 6.2). We give two applications for this unexpected property. For a proper generalized log Calabi–Yau variety, we show that the non-log canonical locus is connected and intersects every log canonical (lc) center (Theorem 6.3). And we obtain an extension theorem from a union of log canonical centers, in the log canonical case (Theorem 6.4). We expect this extension to play a key role in the characterization of the restriction of log canonical rings to lc centers. In §7 we list some questions that appeared naturally during this work.

## 1. Preliminaries

### 1.1 Directed limits

A *directed family* of abelian groups  $(A_m)_{m \in \mathbb{Z}}$  consists of homomorphisms of abelian groups  $\varphi_{mn}: A_m \rightarrow A_n$ , for  $m \leq n$ , such that  $\varphi_{mm} = \text{id}_{A_m}$  and  $\varphi_{np} \circ \varphi_{mn} = \varphi_{mp}$  for  $m \leq n \leq p$ . The *directed limit*  $\varinjlim_m A_m$  of  $(A_m)_{m \in \mathbb{Z}}$  is defined as the quotient of  $\bigoplus_{m \in \mathbb{Z}} A_m$  modulo the subgroup generated by  $x_m - \varphi_{mn}(x_m)$  for all  $m \leq n$  and  $x_m \in A_m$ . The homomorphisms  $\mu_m: A_m \rightarrow \varinjlim_n A_n$ ,  $a_m \mapsto [a_m]$  are compatible with  $\varphi_{mn}$ , and satisfy the following universal property: if  $B$  is an abelian group and  $f_n: A_n \rightarrow B$  are homomorphisms compatible with  $\varphi_{mn}$ , then there exists a unique homomorphism  $f: \varinjlim_m A_m \rightarrow B$  such that  $f_m = f \circ \mu_m$  for all  $m$ . From the explicit description of the directed limit, the following properties hold:  $\varinjlim_n A_n = \bigcup_m \mu_m(A_m)$ , and  $\text{Ker}(A_m \rightarrow \varinjlim_n A_n) = \bigcup_{m \leq n} \text{Ker}(A_m \rightarrow A_n)$ . In particular, we obtain the following lemma.

LEMMA 1.1. *Let  $(A_m)_{m \in \mathbb{Z}}$  be a directed system of abelian groups.*

- (1) *We have that  $A_m \rightarrow \varinjlim_n A_n$  is injective if and only if  $A_m \rightarrow A_n$  is injective for all  $n \geq m$ .*
- (2) *Let  $(B_m)_{m \in \mathbb{Z}}$  be another directed family of abelian groups, let  $f_m: A_m \rightarrow B_m$  be a sequence of compatible homomorphisms. They induce a homomorphism  $f: \varinjlim_m A_m \rightarrow \varinjlim_m B_m$ . If  $f_m$  is injective for  $m \geq m_0$ , then  $f$  is injective.*

### 1.2 Homomorphisms induced in cohomology

For standard notation and results, see Grothendieck [Gro61, 12.1.7, 12.2.5]. Let  $f: X' \rightarrow X$  and  $\pi: X \rightarrow S$  be morphisms of ringed spaces. Denote  $\pi' = \pi \circ f: X' \rightarrow S$ .

Let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module, and  $\mathcal{F}'$  an  $\mathcal{O}_{X'}$ -module. A homomorphism of  $\mathcal{O}_X$ -modules  $u: \mathcal{F} \rightarrow f_* \mathcal{F}'$  induces functorial homomorphisms of  $\mathcal{O}_S$ -modules

$$R^q u: R^q \pi_* \mathcal{F} \rightarrow R^q \pi'_* (\mathcal{F}') \quad (q \geq 0).$$

Grothendieck–Leray constructed a spectral sequence

$$E_2^{pq} = R^p \pi_* (R^q f_* \mathcal{F}') \implies R^{p+q} \pi'_* (\mathcal{F}').$$

LEMMA 1.2. *The homomorphism  $R^1 \pi_* (f_* \mathcal{F}') \rightarrow R^1 \pi'_* (\mathcal{F}')$ , induced by  $\text{id}: f_* \mathcal{F}' \rightarrow f_* \mathcal{F}'$ , is injective.*

*Proof.* The exact sequence of terms of low degree of the Grothendieck–Leray spectral sequence is

$$0 \rightarrow R^1 \pi_* (f_* \mathcal{F}') \rightarrow R^1 \pi'_* (\mathcal{F}') \rightarrow \pi_* (R^1 f_* \mathcal{F}') \rightarrow R^2 \pi_* (f_* \mathcal{F}') \rightarrow R^2 \pi'_* (\mathcal{F}'),$$

and  $R^1 \pi_* (f_* \mathcal{F}') \rightarrow R^1 \pi'_* (\mathcal{F}')$  is exactly the homomorphism induced by the identity of  $f_* \mathcal{F}'$ .  $\square$

The other maps  $R^p \pi_* (f_* \mathcal{F}') \rightarrow R^p \pi'_* (\mathcal{F}')$  ( $p \geq 2$ ), appearing in the spectral sequence as the edge maps  $E_2^{p,0} \rightarrow H^p$ , may not be injective.

Example 1.3. Let  $f: X \rightarrow Y$  be the blow-up at a point of a proper smooth complex surface  $Y$ , let  $E$  be the exceptional divisor. Then the map

$$H^2(Y, f_* \mathcal{O}_X(K_X + E)) \rightarrow H^2(X, \mathcal{O}_X(K_X + E))$$

is not injective. In particular, the Leray spectral sequence for  $f$  and  $\mathcal{O}_X(K_X + E)$  does not degenerate. Indeed, consider the following commutative diagram.

$$\begin{array}{ccc} H^2(X, \mathcal{O}_X(K_X)) & \xrightarrow{\gamma} & H^2(X, \mathcal{O}_X(K_X + E)) \\ \uparrow \alpha & & \uparrow \delta \\ H^2(Y, f_* \mathcal{O}_X(K_X)) & \xrightarrow{\beta} & H^2(Y, f_* \mathcal{O}_X(K_X + E)) \end{array}$$

We have  $R^i f_* \mathcal{O}_X(K_X) = 0$  for  $i = 1, 2$ . Therefore  $\alpha$  is an isomorphism, from the Leray spectral sequence. The natural map  $f_* \mathcal{O}_X(K_X) \rightarrow f_* \mathcal{O}_X(K_X + E)$  is an isomorphism. Therefore  $\beta$  is an isomorphism. By Serre duality, the dual of  $\gamma$  is the inclusion  $\Gamma(X, \mathcal{O}_X(-E)) \rightarrow \Gamma(X, \mathcal{O}_X)$ . Since  $X$  is proper,  $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$ . Therefore  $\Gamma(X, \mathcal{O}_X(-E)) = 0$ . We obtain  $\gamma^\vee = 0$ . Therefore  $\gamma = 0$ .

Since  $\alpha, \beta$  are isomorphisms and  $\gamma = 0$ , we deduce  $\delta = 0$ . But  $H^2(Y, f_* \mathcal{O}_X(K_X + E))$  is non-zero, being isomorphic to  $H^2(X, \mathcal{O}_X(K_X))$ , which is dual to  $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$ . Therefore  $\delta$  is not injective.

### 1.3 Weil divisors

Let  $X$  be a normal algebraic variety defined over  $k$ , an algebraically closed field. A *prime on  $X$*  is a reduced irreducible cycle of codimension one. An  $\mathbb{R}$ -Weil divisor  $D$  on  $X$  is a formal sum

$$D = \sum_E d_E E,$$

where the sum runs after all primes on  $X$ , and  $d_E$  are real numbers such that  $\{E : d_E \neq 0\}$  has at most finitely many elements. It can be viewed as an  $\mathbb{R}$ -valued function defined on all primes, with finite support. By restricting the values to  $\mathbb{Q}$  or  $\mathbb{Z}$ , we obtain the notion of  $\mathbb{Q}$ -Weil divisor and *Weil divisor*, respectively.

Let  $f \in k(X)$  be a rational function. For a prime  $E$  on  $X$ , let  $t$  be a local parameter at the generic point of  $E$ . We define  $v_E(f)$  as the supremum of all  $m \in \mathbb{Z}$  such that  $ft^{-m}$  is regular

at the generic point of  $E$ . If  $f = 0$ , then  $v_E(f) = +\infty$ . Else,  $v_E(f)$  is a well-defined integer. We have  $v_E(fg) = v_E(f) + v_E(g)$  and  $v_E(f + g) \geq \min(v_E(f), v_E(g))$ .

For non-zero  $f \in k(X)$  define  $(f) = \sum_E v_E(f)E$ , where the sum runs after all primes on  $X$ . The sum has finite support, so  $(f)$  is a Weil divisor. A Weil divisor  $D$  on  $X$  is *linearly trivial*, denoted  $D \sim 0$ , if there exists  $0 \neq f \in k(X)$  such that  $D = (f)$ .

DEFINITION 1.4. Let  $D$  be an  $\mathbb{R}$ -Weil divisor on  $X$ . We call  $D$ :

- $\mathbb{R}$ -linearly trivial, denoted  $D \sim_{\mathbb{R}} 0$ , if there exist finitely many  $r_i \in \mathbb{R}$  and  $0 \neq f_i \in k(X)$  such that  $D = \sum_i r_i(f_i)$ ;
- $\mathbb{Q}$ -linearly trivial, denoted  $D \sim_{\mathbb{Q}} 0$ , if there exist finitely many  $r_i \in \mathbb{Q}$  and  $0 \neq f_i \in k(X)$  such that  $D = \sum_i r_i(f_i)$ .

LEMMA 1.5 [Sho93, p. 97]. Let  $E_1, \dots, E_l$  be distinct prime divisors on  $X$ , and  $D$  a  $\mathbb{Q}$ -Weil divisor on  $X$ . If not empty, the set  $\{(x_1, \dots, x_l) \in \mathbb{R}^l : \sum_{i=1}^l x_i E_i \sim_{\mathbb{R}} D\}$  is an affine subspace of  $\mathbb{R}^l$  defined over  $\mathbb{Q}$ .

*Proof.* Case  $D = 0$ : the set  $V_0 = \{x \in \mathbb{R}^l : \sum_{i=1}^l x_i E_i \sim_{\mathbb{R}} 0\}$  is an  $\mathbb{R}$ -vector subspace of  $\mathbb{R}^l$ . Let  $x \in V_0$ . This means that there exist finitely many non-zero rational functions  $f_\alpha \in k(X)^\times$  and finitely many real numbers  $r_\alpha \in \mathbb{R}$  such that

$$\sum_{i=1}^l x_i E_i = \sum_{\alpha} r_\alpha (f_\alpha).$$

This equality of divisors is equivalent to the system of linear equations

$$\text{mult}_E \left( \sum_{i=1}^l x_i E_i \right) = \sum_{\alpha} r_\alpha \text{mult}_E(f_\alpha),$$

one equation for each prime divisor  $E$  which may appear in the support of  $f_\alpha$ , for some  $\alpha$ . We have  $\text{mult}_E(f_\alpha) \in \mathbb{Z}$ . If we fix the  $f_\alpha$ , this means that the  $r_\alpha$  are the solutions of a linear system defined over  $\mathbb{Q}$ , and the corresponding values  $x$  belong to an  $\mathbb{R}$ -vector subspace of  $\mathbb{R}^l$  defined over  $\mathbb{Q}$ .

The above argument shows that  $V_0$  is a union of vector subspaces defined over  $\mathbb{Q}$ . Let  $v_1, \dots, v_k$  be a basis for  $V_0$  over  $\mathbb{R}$ . Each  $v_a$  belongs to some subspace of  $V_0$  defined over  $\mathbb{Q}$ . That is, there exist  $(w_{ab})_b$  in  $V_0 \cap \mathbb{Q}^l$  such that  $v_a \in \sum_b \mathbb{R} w_{ab}$ . It follows that the elements  $w_{ab} \in V_0 \cap \mathbb{Q}^l$  generate  $V_0$  as an  $\mathbb{R}$ -vector space. Therefore  $V_0$  is defined over  $\mathbb{Q}$ .

Case  $D$  arbitrary: suppose  $V = \{x \in \mathbb{R}^l : \sum_{i=1}^l x_i E_i \sim_{\mathbb{R}} D\}$  is non-empty. Let  $x \in V$ . Then  $\sum_{i=1}^l x_i E_i = D + \sum_{\alpha} r_\alpha (f_\alpha)$  for finitely many  $r_\alpha, f_\alpha$  as above. Since  $D$  has rational coefficients, the same argument used above shows that once the  $f_\alpha$  are fixed, there exists another representation  $\sum_{i=1}^l x'_i E_i = D + \sum_{\alpha} r'_\alpha (f_\alpha)$ , with  $x'_i, r'_\alpha \in \mathbb{Q}$ . In particular,  $x' \in V \cap \mathbb{Q}^l$ . We have  $V = x' + V_0$ . Since  $V_0$  is defined over  $\mathbb{Q}$ , we conclude that  $V$  is an affine subspace of  $\mathbb{R}^l$  defined over  $\mathbb{Q}$ .  $\square$

If  $D \sim_{\mathbb{Q}} 0$ , then  $D$  has rational coefficients. If  $D$  has rational coefficients, then  $D \sim_{\mathbb{Q}} 0$  if and only if  $D \sim_{\mathbb{R}} 0$  (by Lemma 1.5).

Let  $D$  be an  $\mathbb{R}$ -divisor on  $X$ . Denote  $D^{=1} = \sum_{d_E=1} E$ ,  $D^{\neq 1} = \sum_{d_E \neq 1} d_E E$ ,  $D^{<0} = \sum_{d_E < 0} d_E E$ ,  $D^{>0} = \sum_{d_E > 0} d_E E$ . The round up (down) of  $D$  is defined as  $\lceil D \rceil = \sum_E \lceil d_E \rceil E$  ( $\lfloor D \rfloor = \sum_E \lfloor d_E \rfloor E$ ), where for  $x \in \mathbb{R}$  we denote  $\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}$  and  $\lceil x \rceil = \min\{n \in \mathbb{Z} : x \leq n\}$ . The fractional part of  $D$  is defined as  $\{D\} = \sum_E \{d_E\} E$ , where for  $x \in \mathbb{R}$  we denote  $\{x\} = x - \lfloor x \rfloor$ .

DEFINITION 1.6. Let  $D$  be an  $\mathbb{R}$ -Weil divisor on  $X$ . We call  $D$   $\mathbb{R}$ -Cartier ( $\mathbb{Q}$ -Cartier, Cartier) if there exists an open covering  $X = \bigcup_i U_i$  such that  $D|_{U_i} \sim_{\mathbb{R}} 0$  ( $D|_{U_i} \sim_{\mathbb{Q}} 0$ ,  $D|_{U_i} \sim 0$ ) for all  $i$ .

### 1.4 Complements of effective Cartier divisors

LEMMA 1.7. *Let  $D$  be an effective Cartier divisor on a Noetherian scheme  $X$ . Let  $U = X \setminus \text{Supp } D$  and consider the open embedding  $w: U \subseteq X$ . Then:*

- (1)  *$w$  is an affine morphism;*
- (2) *let  $\mathcal{F}$  be a quasi-coherent  $\mathcal{O}_X$ -module. The natural inclusions  $\mathcal{F}(mD) \subset \mathcal{F}(nD)$ , for  $m \leq n$ , form a directed family of  $\mathcal{O}_X$ -modules  $(\mathcal{F}(mD))_{m \in \mathbb{Z}}$ ; and*

$$\varinjlim_m \mathcal{F}(mD) = w_*(\mathcal{F}|_U);$$

- (3) *let  $\pi: X \rightarrow S$  be a morphism and  $\mathcal{F}$  a quasi-coherent  $\mathcal{O}_X$ -module. Then  $\varinjlim_m R^q \pi_* \mathcal{F}(mD) \xrightarrow{\sim} R^q(\pi|_U)_*(\mathcal{F}|_U)$  for all  $q$ .*

*Proof.* Let  $X = \bigcup_\alpha V_\alpha$  be an affine open covering such that  $D = (f_\alpha)_\alpha$ , for non-zero divisors  $f_\alpha \in \Gamma(V_\alpha, \mathcal{O}_{V_\alpha})$  such that  $f_\alpha f_\beta^{-1} \in \Gamma(V_\alpha \cap V_\beta, \mathcal{O}_X^\times)$  for all  $\alpha, \beta$ .

The set  $w^{-1}(V_\alpha) = U \cap V_\alpha = D(f_\alpha)$  is affine, so (1) holds. Statement (2) is local, equivalent to the known property

$$\Gamma(D(f_\alpha), \mathcal{F}) = \Gamma(V_\alpha, \mathcal{F})_{f_\alpha} = \varinjlim_m \Gamma(V_\alpha, \mathcal{F}(mD)) = \Gamma\left(V_\alpha, \varinjlim_m \mathcal{F}(mD)\right).$$

For (3), directed limits commute with cohomology on quasi-compact topological spaces. Therefore

$$\varinjlim_m R^q \pi_* \mathcal{F}(mD) \xrightarrow{\sim} R^q \pi_* \left( \varinjlim_m \mathcal{F}(mD) \right) = R^q \pi_* (w_*(\mathcal{F}|_U)).$$

Since  $w$  is affine, the Leray spectral sequence for  $w$  degenerates to isomorphisms

$$R^q \pi_* (w_*(\mathcal{F}|_U)) \xrightarrow{\sim} R^q(\pi|_U)_*(\mathcal{F}|_U).$$

Therefore (3) holds. □

### 1.5 Convention on algebraic varieties

Throughout this paper, a variety is a reduced scheme of finite type over an algebraically closed field  $k$  of characteristic zero.

### 1.6 Explicit Deligne–Du Bois complex for normal crossing varieties

Let  $X$  be a variety with at most *normal crossing singularities*. That is, for every point  $P \in X$ , there exist  $n \geq 1$ ,  $I \subseteq \{1, \dots, n\}$ , and an isomorphism of complete local  $k$ -algebras

$$\frac{k[[T_1, \dots, T_n]]}{(\prod_{i \in I} T_i)} \xrightarrow{\sim} \hat{\mathcal{O}}_{X,P}.$$

Let  $\pi: \bar{X} \rightarrow X$  be the normalization. For  $p \geq 0$ , define the  $\mathcal{O}_X$ -module  $\tilde{\Omega}_{X/k}^p$  to be the image of the natural map  $\Omega_{X/k}^p \rightarrow \pi_* \Omega_{\bar{X}/k}^p$ . We have induced differentials  $d: \tilde{\Omega}_{X/k}^p \rightarrow \tilde{\Omega}_{X/k}^{p+1}$ , and  $\tilde{\Omega}_{X/k}^\bullet$  becomes a differential complex of  $\mathcal{O}_X$ -modules. We call the hypercohomology group  $\mathbb{H}^r(X, \tilde{\Omega}_{X/k}^\bullet)$  the *rth de Rham cohomology group* of  $X/k$ , and denote it by

$$H_{DR}^r(X/k).$$

If the base field is understood, we usually drop it from notation. Let  $X_\bullet$  be the simplicial algebraic variety induced by  $\pi$  (see [Del74]). Its components are  $X_n = (\bar{X}/X)^{\Delta_n}$ , and the simplicial maps are naturally induced. We have a natural augmentation

$$\epsilon: X_\bullet \rightarrow X.$$

We have  $X_0 = \bar{X}$ ,  $X_1 = X_0 \times_X X_0$ ,  $\epsilon_0 = \pi$  and  $\delta_0, \delta_1: X_1 \rightarrow X_0$  are the natural projections. For  $p \geq 0$ , let  $\Omega_{X_\bullet}^p$  be the simplicial  $\mathcal{O}_{X_\bullet}$ -module with components  $\Omega_{X_n}^p$  ( $n \geq 0$ ). The  $\mathcal{O}_X$ -module  $\epsilon_*(\Omega_{X_\bullet}^p)$  is defined as the kernel of the homomorphism

$$\delta_1^* - \delta_0^*: \epsilon_{0*}\Omega_{X_0}^p \rightarrow \epsilon_{1*}\Omega_{X_1}^p.$$

By [DJ74, Lemme 2],  $\epsilon$  is a smooth resolution, and  $R^i\epsilon_*(\Omega_{X_\bullet}^p) = 0$  for  $i > 0, p \geq 0$ .

LEMMA 1.8. *For every  $p$ ,  $\tilde{\Omega}_X^p = \epsilon_*(\Omega_{X_\bullet}^p)$ .*

*Proof.* Since  $\pi \circ \delta_0 = \pi \circ \delta_1$ , we obtain an inclusion  $\tilde{\Omega}_X^p \subseteq \epsilon_*(\Omega_{X_\bullet}^p)$ . The opposite inclusion may be checked locally, in an étale neighborhood of each point. Therefore we may suppose

$$X : \left( \prod_{i=1}^c z_i = 0 \right) \subset \mathbb{A}^{d+1}.$$

Then  $X$  has  $c$  irreducible components  $X_1, \dots, X_c$ , each of them isomorphic to  $\mathbb{A}^d$ . The normalization  $\bar{X}$  is the disjoint union of the  $X_i$ . Therefore  $\Gamma(X, \epsilon_*(\Omega_{X_\bullet}^p))$  consists of  $c$ -uples  $(\omega_1, \dots, \omega_c)$  where  $\omega_i \in \Gamma(X_i, \Omega_{X_i}^p)$  satisfy the cycle condition  $\omega_i|_{X_i \cap X_j} = \omega_j|_{X_i \cap X_j}$  for every  $i < j$ .

By induction on  $c$ , we show that  $\Gamma(X, \epsilon_*(\Omega_{X_\bullet}^p))$  is the image of the homomorphism  $\Gamma(\mathbb{A}^{d+1}, \Omega_{\mathbb{A}^{d+1}}^p) \rightarrow \Gamma(\bar{X}, \Omega_{\bar{X}}^p)$ . The case  $c = 1$  is clear. Suppose  $c \geq 2$ . Let  $\alpha = (\omega_1, \dots, \omega_c)$  be an element of  $\Gamma(X, \epsilon_*(\Omega_{X_\bullet}^p))$ . There exists  $\omega \in \Gamma(\mathbb{A}^{d+1}, \Omega_{\mathbb{A}^{d+1}}^p)$  such that  $\omega_c = \omega|_{X_c}$ . Then we may replace  $\alpha$  by  $\alpha - \omega|_X$ , so that

$$\alpha = (\omega_1, \dots, \omega_{c-1}, 0).$$

The cycle conditions for pairs  $i < c$  give  $\omega_i = z_c \eta_i$ , for some  $\eta_i \in \Gamma(X_i, \Omega_{X_i}^p)$ . The other cycle conditions are equivalent to the fact that  $(\eta_1, \dots, \eta_{c-1}) \in \Gamma(X', \epsilon_*(\Omega_{X'_\bullet}^p))$ , where  $X' : (\prod_{i=1}^{c-1} z_i = 0) \subset \mathbb{A}^{d+1}$ . By induction, there exists  $\eta \in \Gamma(\mathbb{A}^{d+1}, \Omega_{\mathbb{A}^{d+1}}^p)$  such that  $\eta_i = \eta|_{X_i}$  for  $1 \leq i \leq c-1$ . Then  $\alpha = z_c \eta|_X$ .

The map  $\Gamma(\mathbb{A}^{d+1}, \Omega_{\mathbb{A}^{d+1}}^p) \rightarrow \Gamma(\bar{X}, \Omega_{\bar{X}}^p)$  factors through the surjection  $\Gamma(\mathbb{A}^{d+1}, \Omega_{\mathbb{A}^{d+1}}^p) \rightarrow \Gamma(X, \Omega_X^p)$ . Therefore its image is the same as the image of  $\Gamma(X, \Omega_X^p) \rightarrow \Gamma(\bar{X}, \Omega_{\bar{X}}^p)$ .  $\square$

It follows that  $\tilde{\Omega}_X^\bullet \rightarrow R\epsilon_*(\Omega_{X_\bullet}^\bullet)$  is a quasi-isomorphism. From [Del74, Gro66] (see [DuB81, Théorème 4.5]), we deduce the following result.

THEOREM 1.9. *The filtered complex  $(\tilde{\Omega}_X^\bullet, F)$ , where  $F$  is the naive filtration, induces a spectral sequence in hypercohomology*

$$E_1^{pq} = H^q(X, \tilde{\Omega}_X^p) \implies \mathbb{H}^{p+q}(X, \tilde{\Omega}_X^\bullet) = H_{DR}^{p+q}(X/k).$$

*If  $X$  is proper, this spectral sequence degenerates at  $E_1$ .*

Note  $\tilde{\Omega}_X^0 = \mathcal{O}_X$ . If  $d = \dim X$ , then  $\tilde{\Omega}_X^d = \pi_*\Omega_X^d$ , which is a locally free  $\mathcal{O}_X$ -module if and only if  $X$  has no singularities. If  $X$  is non-singular, the natural surjections  $\Omega_X^p \rightarrow \tilde{\Omega}_X^p$  are isomorphisms, for all  $p$ . So our definition of de Rham cohomology for varieties with at most normal crossing singularities is consistent with Grothendieck's definition [Gro66] for non-singular varieties.



### 1.7 Differential forms with logarithmic poles

Let  $(X, \Sigma)$  be a *log smooth pair*, that is  $X$  is a non-singular variety and  $\Sigma$  is an effective divisor with at most normal crossing singularities. Denote  $U = X \setminus \Sigma$ . Let  $w: U \rightarrow X$  be the inclusion. Then  $w_*(\Omega_U^\bullet)$  is the complex of rational differentials on  $X$  which are regular on  $U$ . We identify it with the union of  $\Omega_X^\bullet \otimes \mathcal{O}_X(m\Sigma)$ , after all  $m \geq 0$ .

Let  $p \geq 0$ . The *sheaf of germs of differential  $p$ -forms on  $X$  with at most logarithmic poles along  $\Sigma$* , denoted  $\Omega_X^p(\log \Sigma)$  (see [Del69]), is the sheaf whose sections on an open subset  $V$  of  $X$  are

$$\Gamma(V, \Omega_X^p(\log \Sigma)) = \{\omega \in \Gamma(V, \Omega_X^p \otimes \mathcal{O}_X(\Sigma)) : d\omega \in \Gamma(V, \Omega_X^{p+1} \otimes \mathcal{O}_X(\Sigma))\}.$$

It follows that  $\{\Omega_X^p(\log \Sigma), d^p\}_p$  becomes a subcomplex of  $w_*(\Omega_U^\bullet)$ . It is called the *logarithmic de Rham complex of  $(X, \Sigma)$* , denoted by  $\Omega_X^\bullet(\log \Sigma)$ .

Let  $n = \dim X$ . Then  $\Omega_X^p(\log \Sigma) = 0$  if  $p \notin [0, n]$ . And  $\Omega_X^n(\log \Sigma) = \Omega_X^n \otimes \mathcal{O}_X(\Sigma) = \mathcal{O}_X(K_X + \Sigma)$ , where  $K_X$  is the canonical divisor of  $X$ .

LEMMA 1.10. *Let  $0 \leq p \leq n$ . Then  $\Omega_X^p(\log \Sigma)$  is a coherent locally free extension of  $\Omega_U^p$  to  $X$ . Moreover,  $\Omega_X^0(\log \Sigma) = \mathcal{O}_X$ ,  $\wedge^p \Omega_X^1(\log \Sigma) = \Omega_X^p(\log \Sigma)$ , and the wedge product induces a perfect pairing*

$$\Omega_X^p(\log \Sigma) \otimes_{\mathcal{O}_X} \Omega_X^{n-p}(\log \Sigma) \rightarrow \Omega_X^n(\log \Sigma).$$

*Proof.* The  $\mathcal{O}_X$ -module  $\Omega_X^p(\log \Sigma)$  is coherent, being a subsheaf of  $\Omega_X^p \otimes \mathcal{O}_X(\Sigma)$ . The statements may be checked near a fixed point, after passing to completion. Therefore it suffices to verify the statements at the point  $P = 0$  for  $X = \mathbb{A}_k^n$  and  $\Sigma = (\prod_{i \in J} z_i)$ . As in [EV92, Properties 2.2] for example, it can be checked that in this case  $\Omega_X^p(\log \Sigma)_P$  is the free  $\mathcal{O}_{X,P}$ -module with basis

$$\left\{ \frac{dz^I}{\prod_{i \in J \cap I} z_i} : I \subseteq \{1, \dots, n\}, |I| = p \right\},$$

where for  $I = \{i_1 < \dots < i_p\}$ ,  $dz^I$  denotes  $dz_{i_1} \wedge \dots \wedge dz_{i_p}$ . And  $\prod_{i \in \emptyset} z_i = 1$ . All the statements follow in this case.  $\square$

THEOREM 1.11 [AH55, Del69, EV92, Gro66]. *The inclusion  $\Omega_X^\bullet(\log \Sigma) \subset w_*(\Omega_U^\bullet)$  is a quasi-isomorphism.*

*Proof.* We claim that  $\Omega_X^\bullet(\log \Sigma) \otimes \mathcal{O}_X(D)$  is a subcomplex of  $w_*(\Omega_U^\bullet)$ , for every divisor  $D$  supported by  $\Sigma$ . Indeed, the sheaves in question are locally free, so it suffices to check the statement over the open subset  $X \setminus \text{Sing } \Sigma$ , whose complement has codimension at least two in  $X$ . Therefore we may suppose  $\Sigma$  is non-singular. After passing to completion at a fixed point, it suffices to check the claim at  $P = 0$  for  $X = \mathbb{A}_k^1$  and  $\Sigma = (z)$ . This follows from the formula

$$d(1 \otimes z^m) = m \cdot \frac{dz}{z} \otimes z^m \quad (m \in \mathbb{Z}).$$

We obtain an increasing filtration of  $w_*(\Omega_U^\bullet)$  by sub-complexes

$$\mathcal{K}_m = \Omega_X^*(\log \Sigma) \otimes \mathcal{O}_X(m\Sigma) \quad (m \geq 0).$$

We claim that the quotient complex  $\mathcal{K}_m/\mathcal{K}_{m-1}$  is acyclic, for every  $m > 0$ . Since  $\mathcal{K}_0 = \Omega_X^\bullet(\log \Sigma)$  and  $\bigcup_{m \geq 0} \mathcal{K}_m = w_*(\Omega_U^\bullet)$ , this implies that the quotient complex  $w_*(\Omega_U^\bullet)/\Omega_X^\bullet(\log \Sigma)$  is acyclic, or equivalently  $\Omega_X^\bullet(\log \Sigma) \subset w_*(\Omega_U^\bullet)$  is a quasi-isomorphism.

To prove that  $\mathcal{K}_m/\mathcal{K}_{m-1}$  ( $m > 0$ ) is acyclic, note that we may work locally near a fixed point, and we may also pass to completion (since the components of the two complexes are coherent).

Therefore it suffices to verify the claim at  $P = 0$  for  $X = \mathbb{A}_k^n$  and  $\Sigma = (\prod_{i \in J} z_i)$ . If we denote  $H_j = (z_j)$ , the claim in this case follows from the stronger statement of [EV92, Lemma 2.10]: the inclusion  $\Omega_X^*(\log \Sigma) \otimes \mathcal{O}_X(D) \subset \Omega_X^*(\log \Sigma) \otimes \mathcal{O}_X(D + H_j)$  is a quasi-isomorphism, for every effective divisor  $D$  supported by  $\Sigma$  and every  $j \in J$ .  $\square$

**THEOREM 1.12** [Del71]. *The filtered complex  $(\Omega_X^\bullet(\log \Sigma), F)$ , where  $F$  is the naive filtration, induces a spectral sequence in hypercohomology*

$$E_1^{pq} = H^q(X, \Omega_X^p(\log \Sigma)) \implies \mathbb{H}^{p+q}(X, \Omega_X^\bullet(\log \Sigma)).$$

*If  $X$  is proper, this spectral sequence degenerates at  $E_1$ .*

*Proof.* If  $k = \mathbb{C}$ , the claim follows from [Del71] and GAGA. By the Lefschetz principle, the claim extends to the case when  $k$  is a field of characteristic zero.  $\square$

**LEMMA 1.13.** *For each  $p \geq 0$ , we have a short exact sequence*

$$0 \rightarrow \mathcal{I}_\Sigma \otimes \Omega_X^p(\log \Sigma) \rightarrow \Omega_X^p \rightarrow \tilde{\Omega}_\Sigma^p \rightarrow 0.$$

*Proof.* Let  $\pi: \bar{\Sigma} \rightarrow \Sigma$  be the normalization. We claim that we have an exact sequence

$$0 \rightarrow \mathcal{I}_\Sigma \otimes \Omega_X^p(\log \Sigma) \rightarrow \Omega_X^p \rightarrow \pi_* \Omega_{\bar{\Sigma}}^p,$$

where the second arrow is induced by the inclusion  $\Omega_X^p(\log \Sigma) \subseteq \Omega_X^p \otimes \mathcal{O}_X(\Sigma)$ , and the third arrow is the restriction homomorphism  $\omega \mapsto \omega|_{\bar{\Sigma}}$ . Indeed, denote  $\mathcal{K} = \text{Ker}(\Omega_X^p \rightarrow \pi_* \Omega_{\bar{\Sigma}}^p)$ . We have to show that  $\mathcal{I}_\Sigma \otimes \Omega_X^p(\log \Sigma) = \mathcal{K}$ . This is a local statement which can be checked locally near each point, and since the sheaves are coherent, we may also pass to completion. Therefore it suffices to check the equality at  $P = 0$  in the special case  $X = \mathbb{A}_k^n$ ,  $\Sigma = (\prod_{j \in J} z_j)$ . From the explicit description of local bases for the logarithmic sheaves, the claim holds in this case.

Finally, we compute the image of the restriction. The restriction factors through the *surjection*  $\Omega_X^p \rightarrow \Omega_\Sigma^p$ . Therefore the image coincides with the image of  $\Omega_\Sigma^p \rightarrow \pi_* \Omega_{\bar{\Sigma}}^p$ , which by definition is  $\tilde{\Omega}_\Sigma^p$ .  $\square$

### 1.8 The cyclic covering trick

Let  $X$  be an irreducible normal variety, let  $T$  be a  $\mathbb{Q}$ -Weil divisor on  $X$  such that  $T \sim_{\mathbb{Q}} 0$ . Let  $r \geq 1$  be minimal such that  $rT \sim 0$ . Choose a rational function  $\varphi \in k(X)^\times$  such that  $(\varphi) = rT$ . Denote by

$$\tau': X' \rightarrow X$$

the normalization of  $X$  in the field extension  $k(X) \subseteq k(X)(\sqrt[r]{\varphi})$ . The normal variety  $X'$  is irreducible, since  $r$  is minimal. Choose  $\psi \in k(X')^\times$  such that  $\psi^r = \tau'^* \varphi$ . One computes

$$\tau'_* \mathcal{O}_{X'} = \bigoplus_{i=0}^{r-1} \mathcal{O}_X([iT]) \psi^i.$$

The finite morphism  $\tau'$  is Galois, with Galois group cyclic of order  $r$ . Moreover,  $\tau'$  is étale over  $X \setminus \text{Supp}\{T\}$ .

Suppose now that  $(X, \Sigma)$  is a log smooth pair structure on  $X$ , and the fractional part  $\{T\}$  is supported by  $\Sigma$ . Then  $\tau'$  is flat,  $X'$  has at most quotient singularities (in the étale topology), and  $X' \setminus \tau'^{-1}\Sigma$  is non-singular. Let  $\mu: Y \rightarrow X'$  be an embedded resolution of singularities

of  $(X', \tau'^{-1}\Sigma)$ . If we denote  $\tau = \tau' \circ \mu$ , then  $\tau^{-1}(\Sigma) = \Sigma_Y$  is a normal crossings divisor and  $\mu: Y \setminus \Sigma_Y \rightarrow X' \setminus \tau'^{-1}\Sigma$  is an isomorphism. We obtain the following commutative diagram.

$$\begin{array}{ccc} X' & \xleftarrow{\mu} & Y \\ \tau' \downarrow & \swarrow \tau & \\ X & & \end{array}$$

LEMMA 1.14 [Amb13, EV82]. We have  $R^q \tau_* \Omega_Y^p(\log \Sigma_Y) = 0$  for  $q > 0$ , and

$$\begin{aligned} \tau_* \Omega_Y^p(\log \Sigma_Y) &= \Omega_X^p(\log \Sigma_X) \otimes \tau'_* \mathcal{O}_{X'} \\ &\simeq \bigoplus_{i=0}^{r-1} \Omega_X^p(\log \Sigma_X) \otimes \mathcal{O}_X([iT]). \end{aligned}$$

This statement is proved in [EV82, Lemme 1.2, 1.3] with two extra assumptions:  $X$  is projective, and  $\Sigma$  is a *simple* normal crossing divisor, that is it has normal crossing singularities and its irreducible components are smooth. One can show that the projectivity assumption is not necessary, and the normal crossings case reduces to the simple normal crossing case, by étale base change (see [Amb13]).

## 2. Injectivity for open embeddings

Let  $(X, \Sigma)$  be a log smooth pair, with  $X$  proper. Denote  $U = X \setminus \Sigma$ .

THEOREM 2.1. The restriction homomorphism  $H^q(X, \mathcal{O}_X(K_X + \Sigma)) \rightarrow H^q(U, \mathcal{O}_U(K_U))$  is injective, for all  $q$ .

*Proof.* Consider the inclusion of filtered differential complexes of  $\mathcal{O}_X$ -modules

$$(\Omega_X^\bullet(\log \Sigma), F) \subset (w_*(\Omega_U^\bullet), F),$$

where  $F$  is the naive filtration of a complex. Let  $n = \dim X$ . The inclusion  $F^n \subseteq F^0$  induces the following commutative diagram.

$$\begin{array}{ccc} \mathbb{H}^{q+n}(X, F^n \Omega_X^\bullet(\log \Sigma)) & \xrightarrow{\beta} & \mathbb{H}^{q+n}(X, \Omega_X^\bullet(\log \Sigma)) \\ \alpha^n \downarrow & & \downarrow \alpha \\ \mathbb{H}^{q+n}(X, F^n w_*(\Omega_U^\bullet)) & \longrightarrow & \mathbb{H}^{q+n}(X, w_*(\Omega_U^\bullet)) \end{array}$$

By Theorem 1.11,  $\alpha$  is an isomorphism. Theorem 1.12 implies that  $\beta$  is injective. Therefore  $\alpha \circ \beta$  is injective. Therefore  $\alpha^n$  is injective.

But  $F^n \Omega_X^\bullet(\log \Sigma) = \Omega_X^n(\log \Sigma)[-n]$  and  $F^n w_*(\Omega_U^\bullet) = w_*(\Omega_U^n)[-n]$ . Therefore  $\alpha^n$  becomes

$$\alpha^n: H^q(X, \Omega_X^n(\log \Sigma)) \rightarrow H^q(X, w_*(\Omega_U^n)).$$

The morphism  $w: U \subset X$  is affine, so  $H^q(X, w_*(\Omega_U^n)) \rightarrow H^q(U, \Omega_U^n)$  is an isomorphism. Therefore  $\alpha^n$  becomes the restriction map

$$\alpha^n: H^q(X, \Omega_X^n(\log \Sigma)) \rightarrow H^q(U, \Omega_U^n).$$

□

COROLLARY 2.2. *Let  $T$  be a  $\mathbb{Q}$ -divisor on  $X$  such that  $T \sim_{\mathbb{Q}} 0$  and  $\text{Supp}\{T\} \subseteq \Sigma$ . In particular,  $T|_U$  has integer coefficients. Then the restriction homomorphism*

$$H^q(X, \mathcal{O}_X(K_X + \Sigma + \lfloor T \rfloor)) \rightarrow H^q(U, \mathcal{O}_U(K_U + T|_U))$$

*is injective, for all  $q$ .*

*Proof.* We use the notation of § 1.8. Denote  $V = \tau^{-1}(U) = Y \setminus \Sigma_Y$ . By Theorem 2.1, the restriction

$$H^q(Y, \mathcal{O}_Y(K_Y + \Sigma_Y)) \rightarrow H^q(V, \mathcal{O}_V(K_V))$$

is injective. By the Leray spectral sequence and Lemma 1.14, the restriction

$$H^q(X, \tau_* \mathcal{O}_Y(K_Y + \Sigma_Y)) \rightarrow H^q(U, \tau_* \mathcal{O}_V(K_V))$$

is injective. Equivalently, the direct sum of restrictions

$$\bigoplus_{i=0}^{r-1} (H^q(X, \mathcal{O}_X(K_X + \Sigma + \lfloor iT \rfloor)) \rightarrow H^q(U, \mathcal{O}_U(K_U + iT|_U)))$$

is injective. For  $i = 1$ , we obtain the claim.  $\square$

THEOREM 2.3. *Let  $X$  be a proper non-singular variety. Let  $U$  be an open subset of  $X$  such that  $X \setminus U$  is a normal crossings divisor with irreducible components  $(E_i)_i$ . Let  $L$  be a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{R}} K_X + \sum_i b_i E_i$ , with  $0 < b_i \leq 1$  for all  $i$ . Then the restriction homomorphism*

$$H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(U, \mathcal{O}_U(L|_U))$$

*is injective, for all  $q$ .*

*Proof.* Choose a labeling of the components, say  $E_1, \dots, E_l$ . Since  $L - K_X$  has integer coefficients, it follows by Lemma 1.5 that the set

$$V = \left\{ x \in \mathbb{R}^l : L \sim_{\mathbb{R}} K_X + \sum_{i=1}^l x_i E_i \right\}$$

is a non-empty affine linear subspace of  $\mathbb{R}^l$  defined over  $\mathbb{Q}$ . Then  $(b_1, \dots, b_l) \in V \cap (0, 1]^l$  can be approximated by  $(b'_1, \dots, b'_l) \in V \cap (0, 1]^l \cap \mathbb{Q}^l$ , such that  $b'_i = b_i$  if  $b_i \in \mathbb{Q}$ . Because  $0 \sim_{\mathbb{R}} -L + K_X + \sum_i b'_i E_i$  and the right-hand side has rational coefficients, it follows that  $0 \sim_{\mathbb{Q}} -L + K_X + \sum_i b'_i E_i$ .

In conclusion,  $L \sim_{\mathbb{Q}} K_X + \sum_i b'_i E_i$  and  $0 < b'_i \leq 1$  for all  $i$ . Set  $\Sigma = \sum_i E_i$  and  $T = L - K_X - \sum_i b'_i E_i$ . Then  $T \sim_{\mathbb{Q}} 0$ ,  $\{T\} = \sum_i \{-b'_i\} E_i$  and  $L = K_X + \Sigma + \lfloor T \rfloor$ . Corollary 2.2 gives the claim.  $\square$

REMARK 2.4. Let  $U \subseteq U' \subseteq X$  be another open subset. From the commutative diagram

$$\begin{array}{ccc} H^q(X, \mathcal{O}_X(L)) & \xrightarrow{\quad\quad\quad} & H^q(U, \mathcal{O}_U(L|_U)) \\ & \searrow & \nearrow \\ & H^q(U', \mathcal{O}_{U'}(L|_{U'})) & \end{array}$$

it follows that  $H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(U', \mathcal{O}_{U'}(L|_{U'}))$  is injective for all  $q$ .

*Remark 2.5.* Recall that for an  $\mathcal{O}_X$ -module  $\mathcal{F}$ ,  $\Gamma_\Sigma(X, \mathcal{F})$  is defined as the kernel of  $\Gamma(X, \mathcal{F}) \rightarrow \Gamma(U, \mathcal{F}|_U)$ . The functor  $\Gamma_\Sigma(X, \cdot)$  is left exact. Its derived functors, denoted  $(H_\Sigma^i(X, \mathcal{F}))_{i \geq 0}$ , are called the cohomology of  $X$  modulo  $U$ , with coefficients in  $\mathcal{F}$ . For every  $\mathcal{F}$  we have long exact sequences

$$0 \rightarrow \Gamma_\Sigma(X, \mathcal{F}) \rightarrow \Gamma(X, \mathcal{F}) \rightarrow \Gamma(U, \mathcal{F}|_U) \rightarrow H_\Sigma^1(X, \mathcal{F}) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(U, \mathcal{F}|_U) \rightarrow \cdots$$

Therefore Theorem 2.3 says that the homomorphism  $H_\Sigma^q(X, \mathcal{O}_X(L)) \rightarrow H^q(X, \mathcal{O}_X(L))$  is zero for all  $q$ . Equivalently,  $\Gamma_\Sigma(X, \mathcal{O}_X(L)) = 0$ , and for all  $q$  we have short exact sequences

$$0 \rightarrow H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(U, \mathcal{O}_U(L|_U)) \rightarrow H_\Sigma^{q+1}(X, \mathcal{O}_X(L)) \rightarrow 0.$$

*Remark 2.6.* Theorem 2.3 is also equivalent to the following statement, which generalizes the original result of Esnault and Viehweg [EV92, Theorem 5.1]: let  $D$  be an effective Cartier divisor supported by  $\Sigma$ . Then the long exact sequence induced in cohomology by the short exact sequence  $0 \rightarrow \mathcal{O}_X(L) \rightarrow \mathcal{O}_X(L + D) \rightarrow \mathcal{O}_D(L + D) \rightarrow 0$  breaks up into short exact sequences

$$0 \rightarrow H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(X, \mathcal{O}_X(L + D)) \rightarrow H^q(D, \mathcal{O}_D(L + D)) \rightarrow 0 \quad (q \geq 0).$$

Indeed, let  $D$  be as above. We have the following commutative diagram.

$$\begin{array}{ccc} H^q(X, \mathcal{O}_X(L)) & \xrightarrow{\alpha} & H^q(X, \mathcal{O}_X(L + D)) \\ \downarrow \beta & & \downarrow \\ H^q(U, \mathcal{O}_U(L|_U)) & \xrightarrow{\gamma} & H^q(U, \mathcal{O}_U((L + D)|_U)) \end{array}$$

Since  $D$  is disjoint from  $U$ ,  $\gamma$  is an isomorphism. By Theorem 2.3,  $\beta$  is injective. Therefore  $\gamma \circ \beta$  is injective. It follows that  $\alpha$  is injective. Conversely, suppose  $H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(X, \mathcal{O}_X(L + D))$  is injective for all divisors  $D$  supported by  $X \setminus U$ . Then we see that  $H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(X, \mathcal{O}_X(L + m\Sigma))$  is injective for every  $m \geq 0$ . Lemma 1.1 implies the injectivity of

$$H^q(X, \mathcal{O}_X(L)) \rightarrow \varinjlim_m H^q(X, \mathcal{O}_X(L + m\Sigma)).$$

By Lemma 1.7, this is isomorphic to the homomorphism  $H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(U, \mathcal{O}_U(L|_U))$ .

**COROLLARY 2.7.** *Let  $D$  be an effective Cartier divisor supported by  $\Sigma$ . Then*

$$0 \rightarrow H^q(X, \mathcal{O}_X(K_X + \Sigma)) \rightarrow H^q(X, \mathcal{O}_X(K_X + \Sigma + D)) \rightarrow H^q(D, \mathcal{O}_D(K_X + \Sigma + D)) \rightarrow 0$$

*is a short exact sequence, for all  $q$ .*

*Proof.* By Remark 2.6 for  $L = K_X + \Sigma$ . □

**COROLLARY 2.8.** *The homomorphism  $\Gamma(X, \mathcal{O}_X(K_X + 2\Sigma)) \rightarrow \Gamma(\Sigma, \mathcal{O}_\Sigma(K_X + 2\Sigma))$  is surjective.*

If  $\Sigma$  is the general member of a base point free linear system, this is the original result of Tankeev [Tan71, Proposition 1].

### 3. Differential forms of intermediate degree

Let  $(X, \Sigma)$  be a log smooth pair such that  $X$  is proper and  $U = X \setminus \Sigma$  is contained in an affine open subset of  $X$ .

**THEOREM 3.1.** *We have  $H^q(X, \Omega_X^p(\log \Sigma)) = 0$  for  $p + q > \dim X$ . In particular,  $H^q(X, \mathcal{O}_X(K_X + \Sigma)) = 0$  for  $q > 0$ .*

*Proof.* Consider the logarithmic de Rham complex  $\Omega_X^\bullet(\log \Sigma)$ . Let  $U'$  be an affine open subset of  $X$  containing  $U$ . The inclusions  $U \subseteq U' \subset X$  induce the following commutative diagram.

$$\begin{array}{ccc} \mathbb{H}^r(X, \Omega_X^\bullet(\log \Sigma)) & \xrightarrow{\quad} & \mathbb{H}^r(U, \Omega_U^\bullet) \\ & \searrow & \nearrow \\ & \mathbb{H}^r(U', \Omega_X^\bullet(\log \Sigma)|_{U'}) & \end{array}$$

Since  $U'$  is affine,  $H^q(U', \Omega_X^p(\log \Sigma)|_{U'}) = 0$  for  $q > 0$ . Therefore  $\mathbb{H}^r(U', \Omega_X^\bullet(\log \Sigma)|_{U'})$  is the  $r$ th homology of the differential complex  $\Gamma(U', \Omega_X^\bullet(\log \Sigma))$ . Since  $\Omega_X^p(\log \Sigma) = 0$  for  $p > \dim X$ , we obtain

$$\mathbb{H}^r(U', \Omega_X^\bullet(\log \Sigma)|_{U'}) = 0 \quad \text{for } r > \dim X.$$

Let  $r > \dim X$ . It follows that the horizontal map is zero. But it is an isomorphism by Theorem 1.11. Therefore

$$\mathbb{H}^r(X, \Omega_X^\bullet(\log \Sigma)) = 0.$$

By Theorem 1.12, we have a non-canonical isomorphism

$$\mathbb{H}^r(X, \Omega_X^\bullet(\log \Sigma)) \simeq \bigoplus_{p+q=r} H^q(X, \Omega_X^p(\log \Sigma)).$$

Therefore  $H^q(X, \Omega_X^p(\log \Sigma)) = 0$  for all  $p + q = r$ .  $\square$

Let  $T$  be a  $\mathbb{Q}$ -divisor on  $X$  such that  $T \sim_{\mathbb{Q}} 0$  and  $\text{Supp}\{T\} \subseteq \Sigma$ . In particular,  $T|_U$  has integer coefficients.

**THEOREM 3.2.** *We have  $H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X([T])) = 0$  for  $p + q > \dim X$ . In particular,  $H^q(X, \mathcal{O}_X(K_X + \Sigma + [T])) = 0$  for  $q > 0$ .*

*Proof.* We use the notation of § 1.8. Let  $X \setminus \Sigma \subseteq U'$ , with  $U'$  an affine open subset of  $X$ . Let  $V' = \tau^{-1}(U')$ . By Lemma 1.14, the Leray spectral sequence associated to  $\tau|_{V'}: V' \rightarrow U'$  and  $\Omega_Y^p(\log \Sigma_Y)|_{V'}$  degenerates into isomorphisms

$$H^q(U', (\tau|_{V'})_* \Omega_Y^p(\log \Sigma_Y)|_{V'}) \xrightarrow{\sim} H^q(V', \Omega_Y^p(\log \Sigma_Y)|_{V'}).$$

Since  $U'$  is affine, the left-hand side is zero for  $q > 0$ . Therefore

$$H^q(V', \Omega_Y^p(\log \Sigma_Y)|_{V'}) = 0 \quad \text{for } q > 0.$$

In particular, the spectral sequence

$$E_1^{pq} = H^q(V', \Omega_Y^p(\log \Sigma_Y)|_{V'}) \implies \mathbb{H}^q(V', \Omega_Y^\bullet(\log \Sigma_Y)|_{V'})$$

degenerates into isomorphisms

$$h^r(\Gamma(V', \Omega_Y^\bullet(\log \Sigma_Y))) \simeq \mathbb{H}^r(V', \Omega_Y^\bullet(\log \Sigma_Y)|_{V'}),$$

where the first term is the  $r$ th homology group of the differential complex  $\Gamma(V', \Omega_Y^\bullet(\log \Sigma_Y))$ . Since  $\Omega_Y^p(\log \Sigma_Y) = 0$  for  $p > \dim Y$ , we obtain

$$\mathbb{H}^r(V', \Omega_Y^\bullet(\log \Sigma_Y)|_{V'}) = 0 \quad \text{for } r > \dim Y.$$

Let  $V = \tau^{-1}(U) = Y \setminus \Sigma_Y$ . The restriction map

$$\mathbb{H}^r(Y, \Omega_Y^\bullet(\log \Sigma_Y)) \rightarrow \mathbb{H}^r(V, \Omega_Y^\bullet(\log \Sigma_Y)|_V)$$

is an isomorphism by Theorem 1.11. It factors through  $\mathbb{H}^r(V', \Omega_Y^\bullet(\log \Sigma_Y)|_{V'})$ , hence it is zero for  $r > \dim Y$ . Therefore

$$\mathbb{H}^r(Y, \Omega_Y^\bullet(\log \Sigma_Y)) = 0 \quad \text{for } r > \dim Y.$$

By Theorem 1.12,  $\mathbb{H}^r(Y, \Omega_Y^\bullet(\log \Sigma_Y)) \simeq \bigoplus_{p+q=r} H^q(Y, \Omega_Y^p(\log \Sigma_Y))$ . Therefore

$$H^q(Y, \Omega_Y^p(\log \Sigma_Y)) = 0 \quad \text{for } p + q > \dim Y.$$

The cyclic group of order  $r$  acts on  $H^q(Y, \Omega_Y^p(\log \Sigma_Y))$ , with eigenspace decomposition

$$\bigoplus_{i=0}^{r-1} H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X(\lfloor iT \rfloor)).$$

Therefore  $H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X(\lfloor T \rfloor)) = 0$ .  $\square$

### 3.1 Applications

**COROLLARY 3.3.** *We have  $H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X(-\Sigma - \lfloor T \rfloor)) = 0$  for  $p + q < \dim X$ . In particular,  $H^q(X, \mathcal{O}_X(-\Sigma - \lfloor T \rfloor)) = 0$  for all  $q < \dim X$ .*

*Proof.* This is the dual form of Theorem 3.2, using Serre duality and the isomorphism  $(\Omega_X^p(\log \Sigma))^\vee \simeq \Omega_X^{\dim X - p}(\log \Sigma) \otimes \mathcal{O}_X(-K_X - \Sigma)$ .  $\square$

For  $T = 0$ , we obtain  $H^q(X, \mathcal{I}_\Sigma \otimes \Omega_X^p(\log \Sigma)) = 0$  for all  $p + q < \dim X$ . In particular,  $H^q(X, \mathcal{I}_\Sigma) = 0$  for all  $q < \dim X$ .

**COROLLARY 3.4.** *The homomorphism  $H^q(X, \Omega_X^p \otimes \mathcal{O}_X(-\lfloor T \rfloor)) \rightarrow H^q(\Sigma, \tilde{\Omega}_\Sigma^p \otimes \mathcal{O}_\Sigma(-\lfloor T \rfloor))$  is bijective for  $p + q < \dim \Sigma$  and injective for  $p + q = \dim \Sigma$ .*

*Proof.* Denote  $K^{pq} = H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X(-\Sigma - \lfloor T \rfloor))$ . The short exact sequence of Lemma 1.13 induces a long exact sequence in cohomology

$$K^{pq} \rightarrow H^q(X, \Omega_X^p \otimes \mathcal{O}_X(-\lfloor T \rfloor)) \xrightarrow{\alpha^{qp}} H^q(\Sigma, \tilde{\Omega}_\Sigma^p \otimes \mathcal{O}_\Sigma(-\lfloor T \rfloor)) \rightarrow K^{p, q+1}.$$

By Corollary 3.3,  $\alpha^{qp}$  is bijective for  $q + 1 < \dim X - p$ , and injective for  $q + 1 = \dim X - p$ .  $\square$

**COROLLARY 3.5 (Weak Lefschetz).** *The restriction homomorphism  $H_{DR}^r(X/k) \rightarrow H_{DR}^r(\Sigma/k)$  is bijective for  $r < \dim \Sigma$  and injective for  $r = \dim \Sigma$ .*

*Proof.* Set  $T = 0$ . The homomorphism  $H^q(X, \Omega_X^p) \rightarrow H^q(\Sigma, \tilde{\Omega}_\Sigma^p)$  is bijective for  $p + q < \dim \Sigma$  and injective for  $p + q = \dim \Sigma$ . The Hodge to de Rham spectral sequence degenerates at  $E_1$ , for  $X/k$  by [Del69, Theorem 5.5] and for  $\Sigma/k$  by Theorem 1.9, and is compatible with the maps above.  $\square$

**COROLLARY 3.6.** *Suppose  $\text{Supp}\{T\} = \Sigma$ . Then  $H^q(X, \Omega_X^p(\log \Sigma) \otimes \mathcal{O}_X(\lfloor T \rfloor)) = 0$  for all  $p + q \neq \dim X$ .*

*Proof.* For  $p + q > \dim X$ , this follows from above. For  $p + q < \dim X$ , apply the dual form to  $-T$ , using  $-\Sigma - \lfloor -T \rfloor = \lfloor T \rfloor$ .  $\square$

**COROLLARY 3.7.** *Suppose  $X \setminus \text{Supp}\{T\}$  is contained in an affine open subset of  $X$ . Then  $H^q(X, \mathcal{O}_X([T])) = 0$  for  $q < \dim X$ .*

**THEOREM 3.8** (Akizuki–Nakano). *Let  $X$  be a projective non-singular variety. Let  $L$  be an ample divisor. Then  $H^q(X, \Omega_X^p(L)) = 0$  for  $p+q > \dim X$ . Dually,  $H^q(X, \Omega_X^p(-L)) = 0$  for  $p+q < \dim X$ .*

*Proof.* There exists  $r \geq 1$  such that the general member  $Y \in |rL|$  is non-singular. Set  $T = L - (1/r)Y$  and  $\Sigma = Y$ . Then  $T \sim_{\mathbb{Q}} 0$ ,  $\text{Supp}\{T\} = \Sigma$  and  $X \setminus \Sigma$  is affine. We also have  $[T] = L - Y$ . By Theorem 3.2, we obtain

$$H^q(X, \Omega_X^p(\log Y) \otimes \mathcal{O}_X(L - Y)) = 0 \quad \text{for } p+q > \dim Y.$$

The short exact sequence of Lemma 1.13, tensored by  $L$ , gives an exact sequence

$$H^q(X, \Omega_X^p(\log Y)(L - Y)) \rightarrow H^q(X, \Omega_X^p(L)) \rightarrow H^q(Y, \Omega_Y^p(L)).$$

Let  $p+q > \dim X$ . The first term is zero from above, and the third is zero by induction. Therefore  $H^q(X, \Omega_X^p \otimes \mathcal{O}_X(L)) = 0$ .  $\square$

**COROLLARY 3.9** (Kodaira). *Let  $X$  be a projective non-singular variety. Let  $L$  be an ample divisor on  $X$ . Then  $H^q(X, \mathcal{O}_X(K_X + L)) = 0$  for  $q > 0$ .*

#### 4. Log pairs

A *log pair*  $(X, B)$  consists of a normal algebraic variety  $X$ , endowed with an  $\mathbb{R}$ -Weil divisor  $B$  such that  $K_X + B$  is  $\mathbb{R}$ -Cartier. If  $B$  is effective, we call  $(X, B)$  a *log variety*.

A contraction  $f: X \rightarrow Y$  is a proper morphism such that the natural homomorphism  $\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$  is an isomorphism.

##### 4.1 Totally canonical locus

Let  $(X, B)$  be a log pair. Let  $\mu: X' \rightarrow X$  be a birational contraction such that  $(X', \text{Exc}(\mu) \cup \text{Supp } \mu_*^{-1}B)$  is log smooth. Let

$$\mu^*(K_X + B) = K_{X'} + B_{X'}$$

be the induced log pair structure on  $X'$ . We say that  $\mu: (X', B_{X'}) \rightarrow (X, B)$  is a log crepant birational contraction.

For a prime divisor  $E$  on  $X'$ ,  $1 - \text{mult}_E(B_{X'})$  is called the log discrepancy of  $(X, B)$  in the valuation of  $k(X)$  defined by  $E$ , denoted  $a(E; X, B)$  (see [Amb06] for example).

Define an open subset of  $X$  by the formula  $U = X \setminus \mu(\text{Supp}(B_{X'})^{>0})$ . The definition of  $U$  does not depend on the choice of  $\mu$ , by the following lemma.

**LEMMA 4.1.** *Let  $\mu: (X', B') \rightarrow (X, B)$  be a log crepant proper birational morphism of log pairs with log smooth support. Then  $\mu(\text{Supp } B'^{>0}) = \text{Supp } B^{>0}$ .*

*Proof.* First, we claim that  $B' \leq \mu^*B$ . Indeed,  $X$  is non-singular, so  $K_{X'} - \mu^*K_X$  is effective  $\mu$ -exceptional. From  $\mu^*(K_X + B) = K_{X'} + B'$  we obtain

$$\mu^*B - B' = K_{X'} - \mu^*K_X \geq 0.$$

To prove the statement, denote  $U = X \setminus \text{Supp}(B^{>0})$ . Then  $B|_U \leq 0$ . The claim for  $\mu|_{\mu^{-1}(U)}: (\mu^{-1}(U), B'|_{\mu^{-1}(U)}) \rightarrow (U, B|_U)$  gives  $B'|_{\mu^{-1}(U)} \leq 0$ . Therefore  $\mu(\text{Supp } B'^{>0}) \subseteq \text{Supp } B^{>0}$ . For the opposite inclusion, note that  $\text{Supp } B^{>0}$  has codimension one. Let  $E$  be a prime in  $\text{Supp } B^{>0}$ . Since  $\mu$  is an isomorphism in a neighbourhood of the generic point of  $E$ ,  $E$  also appears as a prime on  $X'$  and  $\text{mult}_E(B') = \text{mult}_E(B) > 0$ . Therefore  $E \subseteq \mu(\text{Supp } B'^{>0})$ .  $\square$



We call  $U$  the *totally canonical locus* of  $(X, B)$ . It is the largest open subset  $U$  of  $X$  with the property that every geometric valuation over  $U$  has log discrepancy at least 1 with respect to  $(U, B|_U)$ . We have

$$X \setminus (\text{Sing}(X) \cup \text{Supp}(B^{>0})) \subseteq U \subseteq X \setminus \text{Supp}(B^{>0}).$$

The first inclusion implies that  $U$  is dense in  $X$ . The second inclusion is an equality if  $(X, \text{Supp } B)$  is log smooth.

#### 4.2 Non-log canonical locus

Let  $(X, B)$  be a log pair with log smooth support. Write  $B = \sum_E b_E E$ , where the sum runs after the prime divisors of  $X$ . Define

$$N(B) = \sum_{b_E < 0} \lfloor b_E \rfloor E + \sum_{b_E > 1} (\lceil b_E \rceil - 1) E.$$

Then  $N(B)$  is a Weil divisor. There exists a unique decomposition  $N(B) = N^+ - N^-$ , where  $N^+, N^-$  are effective divisors with no components in common. Then  $\text{Supp}(N^+) = \text{Supp}(B^{>1})$  and  $\text{Supp}(N^-) = \text{Supp}(B^{<0})$ . We have

$$\lfloor B^{>1} \rfloor - N^+ = \sum_{0 < b_E \in \mathbb{Z}} E.$$

In particular  $N^+ \leq \lfloor B^{>1} \rfloor$ , and the two divisors have the same support. Denote

$$\Delta(B) = B - N(B).$$

We have  $\Delta(B) = \sum_{b_E < 0} \{b_E\} E + \sum_{b_E > 0} (b_E + 1 - \lceil b_E \rceil) E$ . The following properties hold:

- (1) the coefficients of  $\Delta(B)$  belong to the interval  $[0, 1]$ . They are rational if and only if the coefficients of  $B$  are;
- (2)  $\text{Supp}(\Delta(B)) = \text{Supp}(B^{>0}) \cup \bigcup_{0 < b_E \notin \mathbb{Z}} E$ . In particular,  $(X, \Delta(B))$  is a log variety with log canonical singularities and log smooth support;
- (3)  $\text{mult}_E \Delta(B) = 1$  if and only if  $\text{mult}_E B \in \mathbb{Z}_{>0}$ .

LEMMA 4.2. *Let  $\mu: (X', B') \rightarrow (X, B)$  be a log crepant birational contraction of log pairs with log smooth support. Then  $\mu^* N(B) - N(B')$  is an effective  $\mu$ -exceptional divisor. In particular,*

$$\mathcal{O}_X(-N(B)) = \mu_* \mathcal{O}_{X'}(-N(B')).$$

*Proof.* The operation  $B \mapsto N(B)$  is defined componentwise, so  $\mu^* N(B) - N(B')$  is clearly  $\mu$ -exceptional. Decompose  $B = \Delta + N$  and  $B' = \Delta' + N'$ . From  $\mu^*(K + B) = K_{X'} + B'$  we deduce

$$\mu^* N - N' = K_{X'} + \Delta' - \mu^*(K + \Delta).$$

In particular, let  $E$  be a prime divisor on  $X'$ . And  $m_E = \text{mult}_E(\mu^* N - N')$ . Then

$$m_E = a(E; X, \Delta) - a(E; X', \Delta').$$

Since  $(X, \Delta)$  has log canonical singularities and  $\Delta'$  is effective, we obtain

$$m_E \geq 0 - 1 \geq -1.$$

If  $m_E > -1$ , then  $m_E \geq 0$ , as it is an integer. Otherwise,  $m_E = -1$ . In this case  $a(E; X, \Delta) = 0$  and  $a(E; X', \Delta') = 1$ . From  $a(E; X, \Delta) = 0$ , we deduce that  $\mu(E)$  is the transverse intersection

of some components of  $\Delta$  with coefficient 1. That is  $\mu(E)$  is the transverse intersection of some components of  $B$  with coefficients in  $\mathbb{Z}_{\geq 1}$ . In particular,  $B \geq \Delta$  near the generic point of  $\mu(E)$ . We deduce

$$0 = a(E; X, \Delta) \geq a(E; X, B) = a(E; X', B').$$

That is  $\text{mult}_E B' \geq 1$ . Then  $\text{mult}_E \Delta' > 0$ , so  $a(E; X', \Delta') = 1 - \text{mult}_E \Delta' < 1$ . This is a contradiction.  $\square$

DEFINITION 4.3. Let  $(X, B)$  be a log variety. Let  $\mu: (X', B_{X'}) \rightarrow (X, B)$  be a log crepant log resolution. Define

$$\mathcal{I} = \mu_* \mathcal{O}_{X'}(-N(B_{X'})).$$

The coherent  $\mathcal{O}_X$ -module  $\mathcal{I}$  is independent of the choice of  $\mu$ , by Lemma 4.2. Since  $B$  is effective, the divisor  $N(B_{X'})^- = -\lfloor B_{X'}^{\leq 0} \rfloor$  is  $\mu$ -exceptional. Therefore

$$\mathcal{I} \subseteq \mu_* \mathcal{O}_{X'}(N(B_{X'})^-) = \mathcal{O}_X.$$

We call  $\mathcal{I}$  the *ideal sheaf of the non-log canonical locus* of  $(X, B)$ . It defines a closed subscheme  $(X, B)_{-\infty}$  of  $X$  by the short exact sequence

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{(X, B)_{-\infty}} \rightarrow 0.$$

We call  $(X, B)_{-\infty}$  the *locus of non-log canonical singularities* of  $(X, B)$ . It is empty if and only if  $(X, B)$  has log canonical singularities. The complement  $X \setminus (X, B)_{-\infty}$  is the largest open subset on which  $(X, B)$  has log canonical singularities.

Remark 4.4. We introduced in [Amb03] another scheme structure on the locus of non-log canonical singularities of a log variety  $(X, B)$ . The two schemes have the same support, but their structure sheaves usually differ. To compare them, consider a log crepant log resolution  $\mu: (X', B_{X'}) \rightarrow (X, B)$ . Define

$$N^s = \lfloor B_{X'}^{\neq 1} \rfloor = N(B_{X'}) + \sum_{\text{mult}_E(B_{X'}) \in \mathbb{Z}_{>1}} E.$$

Denote  $B_{X'} = \sum_E b_E E$ . Then  $N^s - N(B_{X'}) = \sum_{b_E \in \mathbb{Z}_{>1}} E$  and  $\lfloor B_{X'} \rfloor - N^s = \sum_{b_E=1} E$ . In particular

$$N \leq N^s \leq \lfloor B_{X'} \rfloor.$$

We obtain inclusions of ideal sheaves  $\mu_* \mathcal{O}_{X'}(-N) \supseteq \mu_* \mathcal{O}_{X'}(-N^s) \supseteq \mu_* \mathcal{O}_{X'}(-\lfloor B_{X'} \rfloor)$ . Equivalently, we have closed embeddings of subschemes of  $X$

$$Y \hookrightarrow Y^s \hookrightarrow \text{LCS}(X, B),$$

where  $Y^s$  is the scheme structure introduced in [Amb03] and  $\text{LCS}(X, B)$  is the subscheme structure on the non-klt locus of  $(X, B)$ .

Consider for example the log variety  $(\mathbb{A}^2, 2H_1 + H_2)$ , where  $H_1, H_2$  are the coordinate hyperplanes. The above inclusions are

$$H_1 \hookrightarrow 2H_1 \hookrightarrow 2H_1 + H_2.$$

LEMMA 4.5. Let  $\mu: (X', B') \rightarrow (X, B)$  be a log crepant birational contraction of log pairs with log smooth support. Then  $\mu^* \lfloor B^{\neq 1} \rfloor - \lfloor B_{X'}^{\neq 1} \rfloor$  is an effective  $\mu$ -exceptional divisor. In particular,

$$\mathcal{O}_X(-\lfloor B^{\neq 1} \rfloor) = \mu_* \mathcal{O}_{X'}(-\lfloor B_{X'}^{\neq 1} \rfloor).$$

*Proof.* The operation  $B \mapsto \lfloor B^{\neq 1} \rfloor$  is defined componentwise, so  $\mu^* \lfloor B^{\neq 1} \rfloor - \lfloor B_{X'}^{\neq 1} \rfloor$  is clearly  $\mu$ -exceptional. The equality  $\mu^*(K + B) = K_{X'} + B_{X'}$  becomes

$$\mu^* \lfloor B^{\neq 1} \rfloor - \lfloor B_{X'}^{\neq 1} \rfloor = K_{X'} + B_{X'}^{\neq 1} + \{B_{X'}^{\neq 1}\} - \mu^*(K + B^{\neq 1} + \{B^{\neq 1}\}).$$

Consider the multiplicity of the left-hand side at a prime on  $X'$ . It is an integer. The right-hand side is  $\geq -1$ . If  $> -1$ , it is  $\geq 0$ . Suppose it equals  $-1$ . This implies  $a(E; X, B^{\neq 1} + \{B^{\neq 1}\}) = 0$ . Then  $a(E; X, B^{\neq 1}) = 0$  and  $B = B^{\neq 1}$  near the generic point of  $\mu(E)$ . Then  $a(E; X', B_{X'}) = 0$ . Then the difference is zero. This is a contradiction.  $\square$

### 4.3 Lc centers

For the definition and properties of lc centers, see [Amb06].

LEMMA 4.6. *Let  $(X, B)$  be a log variety with log canonical singularities. Let  $D$  be an effective  $\mathbb{R}$ -Cartier  $\mathbb{R}$ -divisor on  $X$ , let  $Z$  be the union of lc centers of  $(X, B)$  contained in  $\text{Supp } D$ , with reduced structure. Then  $(X, B + \epsilon D)_{-\infty} = Z$  for  $0 < \epsilon \ll 1$ .*

*Proof.* Let  $\mu: X' \rightarrow X$  be a resolution of singularities such that  $(X', \text{Supp } B_{X'} \cup \text{Supp } \mu^* D)$  is log smooth, where  $\mu^*(K_X + B) = K_{X'} + B_{X'}$ , and  $\mu^{-1}(Z)$  has pure codimension one. We have  $\mu^*(K_X + B + \epsilon D) = K_{X'} + B_{X'} + \epsilon \mu^* D$ . Denote

$$\Sigma' = \sum_{\text{mult}_E(B_{X'})=1, \mu(E) \subseteq Z} E.$$

Since the coefficients of  $B_{X'}$  are at most 1, for  $0 < \epsilon \ll 1$  we obtain the formula

$$\begin{aligned} N(B_{X'} + \epsilon \mu^* D) &= \lfloor (B_{X'})^{<0} \rfloor + \sum_{\text{mult}_E(B_{X'})=1, \mu(E) \subseteq \text{Supp } D} E \\ &= \lfloor (B_{X'})^{<0} \rfloor + \Sigma'. \end{aligned}$$

Denote  $A = -\lfloor (B_{X'})^{<0} \rfloor$ , an effective  $\mu$ -exceptional divisor on  $X'$ . Consider the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mu_* \mathcal{O}_{X'}(A - \Sigma') & \longrightarrow & \mu_* \mathcal{O}_{X'}(A) & \xrightarrow{r} & \mu_* \mathcal{O}_{\Sigma'}(A|_{\Sigma'}) \xrightarrow{\partial} R^1 \mu_* \mathcal{O}_{X'}(A - \Sigma') \\ & & \uparrow \alpha & & \uparrow \beta & & \uparrow \gamma \\ 0 & \longrightarrow & \mathcal{I}_Z & \longrightarrow & \mathcal{O}_X & \longrightarrow & \mathcal{O}_Z \longrightarrow 0 \end{array}$$

We claim that  $\partial = 0$ . Indeed, denote  $B' = \{B_{X'}^{<0}\} + B_{X'}^{>0} - \Sigma'$ . Then  $A - \Sigma' \sim_{\mathbb{R}} K_{X'} + B'$  over  $X$ ,  $(X', B')$  has log canonical singularities, and  $\mu(C) \not\subseteq Z$  for every lc center  $C$  of  $(X', B')$ . The sheaf  $\mu_* \mathcal{O}_{\Sigma'}(A|_{\Sigma'})$  is supported by  $Z$ , so the image of  $\partial$  is supported by  $Z$ . Suppose by contradiction that  $\partial$  is non-zero. Let  $s$  be a non-zero local section of  $\text{Im } \partial$ . By [Amb03, Theorem 3.2(i)],  $(X', B')$  admits an lc center  $C$  such that  $\mu(C) \subseteq \text{Supp}(s)$ . Since  $\text{Supp}(s) \subseteq Z$ , we obtain  $\mu(C) \subseteq Z$ , a contradiction.

Since  $A$  is effective and  $\mu$ -exceptional,  $\beta$  is an isomorphism. The map  $\gamma$  is injective. Since  $r$  is surjective,  $\gamma$  is also surjective, hence an isomorphism. We conclude that  $\alpha$  is an isomorphism. That is  $\mathcal{I}_Z = \mu_* \mathcal{O}_{X'}(-N(B_{X'} + \epsilon \mu^* D)) = \mathcal{I}_{(X, B + \epsilon D)_{-\infty}}$ .  $\square$

### 5. Injectivity for log varieties

**THEOREM 5.1.** *Let  $(X, B)$  be a proper log variety with log canonical singularities. Let  $U$  be the totally canonical locus of  $(X, B)$ . Let  $L$  be a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{R}} K + B$ . Then the restriction homomorphism*

$$H^1(X, \mathcal{O}_X(L)) \rightarrow H^1(U, \mathcal{O}_U(L|_U))$$

*is injective.*

*Proof.* Let  $\mu: X' \rightarrow X$  be a birational contraction such that  $X'$  is non-singular, the exceptional locus  $\text{Exc } \mu$  has codimension one, and  $\text{Exc } \mu \cup \text{Supp}(\mu_*^{-1}B)$  has normal crossings. We can write

$$K_{X'} + \mu_*^{-1}B + \text{Exc } \mu = \mu^*(K + B) + A,$$

with  $A$  supported by  $\text{Exc } \mu$ . Since  $(X, B)$  has log canonical singularities,  $A$  is effective. Denote  $B' = \mu_*^{-1}B + \text{Exc } \mu - \{A\}$  and  $L' = \mu^*L + \lfloor A \rfloor$ . We obtain

$$L' \sim_{\mathbb{R}} K_{X'} + B'.$$

Denote  $U' = X' \setminus B'$ . We claim that  $U' \subseteq \mu^{-1}(U)$ . Indeed, this is equivalent to the inclusion

$$\text{Supp}(B') \supseteq \mu^{-1}\mu(\text{Supp } B_{X'}^{>0}).$$

By Zariski's Main Theorem,  $\text{Exc } \mu = \mu^{-1}(X \setminus V)$ , where  $V$  is the largest open subset of  $X$  such that  $\mu$  is an isomorphism over  $V$ . Over  $X \setminus V$ , the inclusion is clear since  $\text{Exc } \mu \subseteq \text{Supp } B'$ . Over  $V$ ,  $\mu$  is an isomorphism and the inclusion becomes an equality. This proves the claim.

Since  $A$  is effective and  $\mu_*A = 0$ , we have  $\mathcal{O}_X(L) \xrightarrow{\sim} \mu_*\mathcal{O}_{X'}(L')$ . From  $U' \subseteq \mu^{-1}(U)$  we obtain the following commutative diagram.

$$\begin{array}{ccc} H^1(X', \mathcal{O}_{X'}(L')) & \xrightarrow{\alpha'} & H^1(U', \mathcal{O}_{U'}(L'|_{U'})) \\ \beta \uparrow & & \uparrow \\ H^1(X, \mathcal{O}_X(L)) & \xrightarrow{\alpha} & H^1(U, \mathcal{O}_U(L|_U)) \end{array}$$

By Theorem 2.3,  $\alpha'$  is injective. Since  $\mathcal{O}_X(L) = \mu_*\mathcal{O}_{X'}(L')$ , Lemma 1.2 implies that  $\beta$  is injective. Then  $\alpha' \circ \beta$  is injective. The diagram is commutative, so  $\alpha$  is injective.  $\square$

**COROLLARY 5.2.** *In the assumptions of Theorem 5.1, let  $D$  be an effective Cartier divisor such that  $\text{Supp}(D) \cap U = \emptyset$ . Then we have a short exact sequence*

$$0 \rightarrow \Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(X, \mathcal{O}_X(L + D)) \rightarrow \Gamma(D, \mathcal{O}_D(L + D)) \rightarrow 0.$$

*Proof.* Consider the following commutative diagram.

$$\begin{array}{ccc} H^1(X, \mathcal{O}_X(L)) & \xrightarrow{\alpha} & H^1(X, \mathcal{O}_X(L + D)) \\ \downarrow \beta & & \downarrow \\ H^1(U, \mathcal{O}_U(L|_U)) & \xrightarrow{\gamma} & H^1(U, \mathcal{O}_U((L + D)|_U)) \end{array}$$

Since  $D$  is disjoint from  $U$ ,  $\gamma$  is an isomorphism. Since  $\beta$  is injective, we obtain that  $\gamma \circ \beta$  is injective. Therefore  $\alpha$  is injective. The long exact sequence induced in cohomology by the short exact sequence  $0 \rightarrow \mathcal{O}_X(L) \rightarrow \mathcal{O}_X(L + D) \rightarrow \mathcal{O}_D(L + D) \rightarrow 0$  gives the claim.  $\square$

### 5.1 Applications

Let  $(X, B)$  be a proper log variety with log canonical singularities, let  $L, H$  be Cartier divisors on  $X$ .

**COROLLARY 5.3.** *Suppose  $L \sim_{\mathbb{R}} K_X + B$ . Suppose the totally canonical locus of  $(X, B)$  is contained in some affine open subset  $U' \subseteq X$ . Then  $H^1(X, \mathcal{O}_X(L)) = 0$ .*

*Proof.* Let  $U$  be the totally canonical locus of  $(X, B)$ . The restriction homomorphism  $H^1(X, \mathcal{O}_X(L)) \rightarrow H^1(U, \mathcal{O}_U(L|_U))$  is injective. It factors through  $H^1(U', \mathcal{O}_{U'}(L|_{U'})) = 0$ , hence it is zero. Therefore  $H^1(X, \mathcal{O}_X(L)) = 0$ .  $\square$

**COROLLARY 5.4.** *Let  $L \sim_{\mathbb{R}} K_X + B$ . Let  $H$  be a Cartier divisor on  $X$  such that the linear system  $|nH|$  is base point free for some positive integer  $n$ . Let  $m_0 \geq 1$  and  $s \in \Gamma(X, \mathcal{O}_X(m_0 H))$  such that  $s|_C \neq 0$  for every lc center of  $(X, B)$ . Then the multiplication*

$$\otimes s: H^1(X, \mathcal{O}_X(L + mH)) \rightarrow H^1(X, \mathcal{O}_X(L + (m + m_0)H))$$

*is injective for  $m \geq 1$ .*

*Proof.* Let  $D$  be the zero locus of  $s$ . There exists a rational number  $0 < \epsilon < 1/m_0$  such that  $(X, B + \epsilon D)$  has log canonical singularities. We have

$$L + mH \sim_{\mathbb{R}} K_X + B + \epsilon D + (m - \epsilon m_0)H.$$

There exists  $n \geq 1$  such that the linear system  $|n(m - \epsilon m_0)H|$  has no base points. Let  $Y$  be a general member, and denote  $B' = B + \epsilon D + (1/n)Y$ . Then  $(X, B')$  has log canonical singularities,  $\text{Supp } D \subseteq \text{Supp } B'$  and

$$L + mH \sim_{\mathbb{R}} K_X + B'.$$

Since  $\text{Supp}(D)$  is disjoint from the totally canonical locus of  $(X, B')$ , Corollary 5.2 gives the injectivity of  $H^1(X, L + mH) \rightarrow H^1(X, L + mH + D)$ .  $\square$

**COROLLARY 5.5.** *Let  $V \subseteq \Gamma(X, \mathcal{O}_X(H))$  be a vector subspace such that  $V \otimes_k \mathcal{O}_X \rightarrow \mathcal{O}_X(H)$  is surjective. If  $L \sim_{\mathbb{R}} K + B + tH$  and  $t > \dim_k V$ , then the multiplication map*

$$V \otimes_k \Gamma(X, \mathcal{O}_X(L - H)) \rightarrow \Gamma(X, \mathcal{O}_X(L))$$

*is surjective.*

*Proof.* We use induction on  $\dim V$ . If  $\dim V = 1$ , then  $V = k\varphi$ , with  $\varphi: \mathcal{O}_X \xrightarrow{\sim} \mathcal{O}_X(H)$ . Then  $\otimes \varphi: \mathcal{O}_X(L - H) \rightarrow \mathcal{O}_X(L)$  is an isomorphism, so the claim holds.

Let  $\dim V > 1$ . Let  $\varphi \in V$  be a general element, let  $Y = (\varphi) + H$ . Then the claim is equivalent to the surjectivity of the homomorphism

$$V|_Y \otimes \Gamma(X, \mathcal{O}_X(L - H))|_Y \rightarrow \Gamma(X, \mathcal{O}_X(L))|_Y$$

where  $\Gamma(X, \mathcal{F})|_Y$  denotes the image of the restriction map  $\Gamma(X, \mathcal{F}) \rightarrow \Gamma(Y, \mathcal{F} \otimes \mathcal{O}_Y)$ , and  $V|_Y$  is the image of  $V$  under this restriction for  $\mathcal{F} = \mathcal{O}_X(H)$ .

Assuming  $\Gamma(X, \mathcal{O}_X(L - H))|_Y = \Gamma(Y, \mathcal{O}_Y(L))$  and  $\Gamma(X, \mathcal{O}_X(L - H))|_Y = \Gamma(Y, \mathcal{O}_Y(L - H))$ , we prove the claim as follows: we have  $L \sim_{\mathbb{R}} K_X + B + Y + (t - 1)H$ . By adjunction, using that  $Y$  is general, we have  $L|_Y \sim_{\mathbb{R}} K_Y + B|_Y + (t - 1)H|_Y$ ,  $(Y, B|_Y)$  has log canonical singularities, and  $t - 1 > \dim V - 1 = \dim V|_Y$ . Therefore  $V|_Y \otimes \Gamma(Y, \mathcal{O}_Y(L - H)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L))$  is surjective by induction.

It remains to show that  $\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L))$  and  $\Gamma(X, \mathcal{O}_X(L - H)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L - H))$  are surjective. Consider the second homomorphism. We have

$$L - Y \sim_{\mathbb{R}} K_X + B + (t - 1)H = K_X + B + \epsilon Y + (t - 1 - \epsilon)H.$$

Since  $Y$  is general,  $(X, B + \epsilon Y)$  has log canonical singularities for  $0 < \epsilon \ll 1$ . Since  $H$  is free, we deduce that  $L - Y \sim_{\mathbb{R}} K_X + B'$  with  $(X, B')$  having log canonical singularities, and  $Y \subseteq \text{Supp } B'$ . By Corollary 5.2,  $\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L))$  is surjective. The surjectivity of the other homomorphism is proved in the same way.  $\square$

## 6. Restriction to the non-log canonical locus

Let  $(X, B)$  be a proper log variety, and  $L$  a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{R}} K_X + B$ . Suppose the locus of non-log canonical singularities  $Y = (X, B)_{-\infty}$  is non-empty.

LEMMA 6.1. *Suppose  $(X, \text{Supp } B)$  is log smooth.*

(1) *The long exact sequence induced in cohomology by the short exact sequence*

$$0 \rightarrow \mathcal{I}_Y(L) \rightarrow \mathcal{O}_X(L) \rightarrow \mathcal{O}_Y(L) \rightarrow 0$$

*breaks up into short exact sequences*

$$0 \rightarrow H^q(X, \mathcal{I}_Y(L)) \rightarrow H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(Y, \mathcal{O}_Y(L)) \rightarrow 0 \quad (q \geq 0).$$

(2) *Let  $E$  be a prime divisor on  $X$  such that  $\text{mult}_E B = 1$ . The long exact sequence induced in cohomology by the short exact sequence*

$$0 \rightarrow \mathcal{I}_Y(L - E) \rightarrow \mathcal{O}_X(L - E) \rightarrow \mathcal{O}_Y(L - E) \rightarrow 0$$

*breaks up into short exact sequences*

$$0 \rightarrow H^q(X, \mathcal{I}_Y(L - E)) \rightarrow H^q(X, \mathcal{O}_X(L - E)) \rightarrow H^q(Y, \mathcal{O}_Y(L - E)) \rightarrow 0 \quad (q \geq 0).$$

*Proof.* (1) Let  $N = N(B)$ , so that  $\mathcal{I}_Y = \mathcal{O}_X(-N)$ . We have  $L - N \sim_{\mathbb{R}} K_X + \Delta$  and  $N$  is supported by  $\Delta$ . By Remark 2.6, the natural map  $H^q(X, \mathcal{O}_X(L - N)) \rightarrow H^q(X, \mathcal{O}_X(L))$  is injective for all  $q$ .

(2) We have  $L - E \sim_{\mathbb{R}} K_X + B - E$  and  $(X, B - E)_{-\infty} = (X, B)_{-\infty} = Y$ . Therefore (2) follows from (1).  $\square$

THEOREM 6.2 (Extension from non-lc locus). *We have a short exact sequence*

$$0 \rightarrow \Gamma(X, \mathcal{I}_Y(L)) \rightarrow \Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Y, \mathcal{O}_Y(L)) \rightarrow 0.$$

*Proof.* Let  $\mu: (X', B_{X'}) \rightarrow (X, B)$  be a log crepant log resolution. Let  $N(B_{X'}) = N = N^+ - N^-$  and  $\Delta = B_{X'} - N(B_{X'})$ . We have

$$\mu^*L - N \sim_{\mathbb{R}} K_{X'} + \Delta$$

and  $N^+$  is supported by  $\Delta$ . By Remark 2.6, we obtain for all  $q$  short exact sequences

$$0 \rightarrow H^q(X', \mathcal{O}_{X'}(\mu^*L - N)) \rightarrow H^q(X', \mathcal{O}_{X'}(\mu^*L + N^-)) \rightarrow H^q(N', \mathcal{O}_{N'}(\mu^*L + N^-)) \rightarrow 0.$$

By definition,  $\mathcal{I}_Y = \mu_* \mathcal{O}_{X'}(-N)$ . Thus  $\mathcal{I}_Y(L) = \mu_* \mathcal{O}_{X'}(\mu^* L - N)$ , and we obtain the following commutative diagram.

$$\begin{array}{ccc} H^q(X', \mathcal{O}_{X'}(\mu^* L + N^- - N^+)) & \xrightarrow{\gamma^q} & H^q(X', \mathcal{O}_{X'}(\mu^* L + N^-)) \\ \beta^q \uparrow & & \uparrow \\ H^q(X, \mathcal{I}_Y(L)) & \xrightarrow{\alpha^q} & H^q(X, \mathcal{O}_X(L)) \end{array}$$

From above,  $\gamma^q$  is injective. By Lemma 1.2,  $\beta^1$  is injective. Therefore  $\gamma^1 \circ \beta^1$  is injective. Therefore  $\alpha^1$  is injective, which is equivalent to our statement.  $\square$

### 6.1 Applications

The first application was first stated by Shokurov, who showed that it follows from the log minimal model program and log abundance in the same dimension (see the proof of [Sho03, Lemma 10.15]).

**THEOREM 6.3** (Global inversion of adjunction). *Let  $(X, B)$  be a proper connected log variety such that  $K_X + B \sim_{\mathbb{R}} 0$ . Suppose  $Y = (X, B)_{-\infty}$  is non-empty. Then  $Y$  is connected, and intersects every lc center of  $(X, B)$ .*

*Proof.* By Theorem 6.2, we have a short exact sequence

$$0 \rightarrow \Gamma(X, \mathcal{I}_Y) \rightarrow \Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(Y, \mathcal{O}_Y) \rightarrow 0.$$

We have  $0 = \Gamma(X, \mathcal{I}_Y), k \xrightarrow{\sim} \Gamma(X, \mathcal{O}_X)$ . Therefore  $k \xrightarrow{\sim} \Gamma(Y, \mathcal{O}_Y)$ , so  $Y$  is connected.

Let  $C$  be a log canonical center of  $(X, B)$ . Let  $\mu: (X', B_{X'}) \rightarrow (X, B)$  be a log resolution such that  $\mu^{-1}(C)$  has codimension one. Let  $\Sigma$  be the part of  $B_{X'}^{-1}$  contained in  $\mu^{-1}(C)$ . We have  $\mu(\Sigma) = C$ . Let  $B' = B_{X'} - \Sigma$  and  $N = N(B') = N(B_{X'})$ . We have

$$-\Sigma - N \sim_{\mathbb{R}} K_{X'} + \Delta(B').$$

The boundary  $\Delta(B')$  supports  $N^+$ . By Remark 2.6, we obtain a surjection

$$\Gamma(X', \mathcal{O}_{X'}(-\Sigma + N^-)) \rightarrow \Gamma(N^+, \mathcal{O}_{N^+}(-\Sigma + N^-)).$$

We have  $\Gamma(X', \mathcal{O}_{X'}(-\Sigma + N^-)) \subseteq \Gamma(X, \mathcal{I}_C) = 0$ . Therefore  $\Gamma(X', \mathcal{O}_{X'}(-\Sigma + N^-)) = 0$ . We obtain  $\Gamma(N^+, \mathcal{O}_{N^+}(-\Sigma + N^-)) = 0$ . Since

$$0 = \Gamma(N^+, \mathcal{O}_{N^+}(-\Sigma + N^-)) \subseteq \Gamma(N^+, \mathcal{O}_{N^+}(N^-)) \neq 0,$$

we infer  $\Sigma \cap N^+ \neq \emptyset$ . This implies  $C \cap Y \neq \emptyset$ .  $\square$

The next application is a corollary of [Amb03, Theorem 4.4.], if  $H$  is  $\mathbb{Q}$ -ample.

**THEOREM 6.4** (Extension from lc centers). *Let  $(X, B)$  be a proper log variety with log canonical singularities. Let  $L$  be a Cartier divisor on  $X$  such that  $H = L - (K_X + B)$  is a semiample  $\mathbb{Q}$ -divisor. Let  $m_0 \geq 1$ ,  $D \in |m_0 H|$ , and denote by  $Z$  the union of lc centers of  $(X, B)$  contained in  $\text{Supp } D$ . Then the restriction homomorphism*

$$\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Z, \mathcal{O}_Z(L))$$

*is surjective.*

*Proof.* By Lemma 4.6, there exists  $\epsilon \in (0, 1) \cap \mathbb{Q}$  such that  $(X, B + \epsilon D)_{-\infty} = Z$ . Let  $m_1 \geq 1$  such that the linear system  $|m_1 H|$  has no base points. Let  $D' \in |m_1 H|$  be a general member. Then  $(X, B + \epsilon D + (1/m_1 - \epsilon/m_0 m_1) D')_{-\infty} = Z$  and

$$L \sim_{\mathbb{Q}} K_X + B + \epsilon D + \left( \frac{1}{m_1} - \frac{\epsilon}{m_0 m_1} \right) D'.$$

By Theorem 6.2,  $\Gamma(X, \mathcal{O}_X(L)) \rightarrow \Gamma(Z, \mathcal{O}_Z(L))$  is surjective.  $\square$

**COROLLARY 6.5.** *Let  $(X, B)$  be a proper log variety with log canonical singularities such that the linear system  $|m_1(K_X + B)|$  has no base points for some  $m_1 \geq 1$ . Let  $m_0 \geq 1$ ,  $D \in |m_0(K_X + B)|$ , and denote by  $Z$  the union of lc centers of  $(X, B)$  contained in  $\text{Supp } D$ . Then*

$$\Gamma(X, \mathcal{O}_X(mK_X + mB)) \rightarrow \Gamma(Z, \mathcal{O}_Z(mK_X + mB))$$

*is surjective for every  $m \geq 2$  such that  $mK_X + mB$  is Cartier.*

*Proof.* Apply Theorem 6.4 to  $m(K_X + B) = K_X + B + (m - 1)(K_X + B)$ .  $\square$

## 7. Questions

*Question 7.1.* Let  $(X, \Sigma)$  be a log smooth pair, with  $X$  proper. Denote  $U = X \setminus \Sigma$ . Is the restriction  $H^q(X, \Omega_X^p(\log \Sigma)) \rightarrow H^q(U, \Omega_U^p)$  injective for  $p + q > \dim X$ ?

*Example 7.2.* Let  $P \in S$  be the germ of non-singular point, of dimension  $d \geq 2$ . Let  $\mu: X \rightarrow S$  be the blow-up at  $P$ , with exceptional locus  $E \simeq \mathbb{P}^{d-1}$ . Denote  $U = X \setminus E$ . The residue map

$$R^{d-1} \mu_* \mathcal{O}_X(K_X + E) \rightarrow R^{d-1} \mu_* \mathcal{O}_E(K_E)$$

is an isomorphism, so  $R^{d-1} \mu_* \mathcal{O}_X(K_X + E)$  is a skyscraper sheaf on  $X$  centered at  $P$ . Since  $\mu$  is an isomorphism on  $U$ ,  $R^{d-1}(\mu|_U)_* \mathcal{O}_E(K_E) = 0$ . Therefore the restriction homomorphism

$$R^{d-1} \mu_* \mathcal{O}_X(K_X + E) \rightarrow R^{d-1}(\mu|_U)_* \mathcal{O}_U(K_U)$$

is not injective.

*Question 7.3.* Let  $(X, \Sigma)$  be a log smooth pair. Denote  $U = X \setminus \Sigma$ . Let  $\pi: X \rightarrow S$  be a proper morphism, let  $\pi|_U: U \rightarrow S$  be its restriction to  $U$ . Suppose that  $\pi(C) = \pi(X)$  for every strata  $C$  of  $(X, \Sigma)$ . Is the restriction  $R^q \pi_* \mathcal{O}_X(K_X + \Sigma) \rightarrow R^q(\pi|_U)_* \mathcal{O}_U(K_U)$  injective for all  $q$ ?

*Question 7.4.* Let  $(X, B)$  be a proper log variety with log canonical singularities. Let  $U$  be the totally canonical locus of  $(X, B)$ . Let  $L$  be a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{R}} K_X + B$ . Is the restriction  $H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(U, \mathcal{O}_U(L|_U))$  injective for all  $q$ ?

*Question 7.5.* Let  $(X, B)$  be a proper log variety. Suppose the locus of non-log canonical singularities  $Y = (X, B)_{-\infty}$  is non-empty. Let  $L$  be a Cartier divisor on  $X$  such that  $L \sim_{\mathbb{R}} K_X + B$ . Does the long exact sequence induced in cohomology by the short exact sequence  $0 \rightarrow \mathcal{I}_Y(L) \rightarrow \mathcal{O}_X(L) \rightarrow \mathcal{O}_Y(L) \rightarrow 0$  break up into short exact sequences

$$0 \rightarrow H^q(X, \mathcal{I}_Y(L)) \rightarrow H^q(X, \mathcal{O}_X(L)) \rightarrow H^q(Y, \mathcal{O}_Y(L)) \rightarrow 0 \quad (q \geq 0)?$$

## ACKNOWLEDGEMENTS

I would like to thank Paolo Cascini, Alexandru Dimca, János Kollár, Adrian Langer, Vyacheslav V. Shokurov and the anonymous referees for useful discussions, comments and corrections.



## REFERENCES

- Amb03 F. Ambro, *Quasi-log varieties*, in *Birational geometry: linear systems and finitely generated algebras: collected papers*, Proceedings of the Steklov Institute of Mathematics, vol. 240, eds V. A. Iskovskikh and V. V. Shokurov (Nauka, Moscow, 2003), 220–239.
- Amb06 F. Ambro, *Basic properties of log canonical centers*. Preprint (2006), [arXiv:math/0611205](https://arxiv.org/abs/math/0611205).
- Amb13 F. Ambro, *Cyclic covers and toroidal embeddings*. Preprint (2013), [arXiv:1310.3951](https://arxiv.org/abs/1310.3951).
- AH55 M. F. Atiyah and W. V. D. Hodge, *Integrals of the second kind on an algebraic variety*, Ann. of Math. (2) **62** (1955), 56–91.
- Del69 P. Deligne, *Théorème de Lefschetz et critères de dégénérescence de suites spectrales*, Publ. Math. Inst. Hautes Études Sci. **35** (1969), 107–126.
- Del P. Deligne, *Equations différentielles à points singulières régulières*, Lecture Notes in Mathematics, vol. 163 (Springer, Heidelberg, 1970).
- Del71 P. Deligne, *Théorie de Hodge II*, Publ. Math. Inst. Hautes Études Sci. **40** (1971), 5–57.
- Del74 P. Deligne, *Théorie de Hodge III*, Publ. Math. Inst. Hautes Études Sci. **44** (1974), 5–77.
- DuB81 P. Du Bois, *Complexe de de Rham filtré d’une variété singulière*, Bull. Soc. Math. France **109** (1981), 41–81.
- DJ74 P. Du Bois and P. Jarraud, *Une propriété de commutation au changement de base des images directes supérieures de faisceaux structural*, C. R. Acad. Sci. Paris, Sér. A **279** (1974), 745–747.
- EV82 H. Esnault and E. Viehweg, *Revêtements cycliques*, in *Algebraic threefolds (Varenna, 1981)*, Lecture Notes in Mathematics, vol. 947 (Springer, Berlin–New York, 1982), 241–250.
- EV86 H. Esnault and E. Viehweg, *Logarithmic de Rham complexes and vanishing theorems*, Invent. Math. **86** (1986), 161–194.
- EV92 H. Esnault and E. Viehweg, *Lectures on vanishing theorems*, in *DMV Seminar* (Birkhäuser, Basel, 1992).
- Gro61 A. Grothendieck, *Eléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): III. Étude cohomologique des faisceaux cohérents, Première partie*, Publ. Math. Inst. Hautes Études Sci. **11** (1961), 5–167.
- Gro66 A. Grothendieck, *On the de Rham cohomology of algebraic varieties*, Publ. Math. Inst. Hautes Études Sci. **29** (1966), 95–103.
- Kaw82 Y. Kawamata, *A generalization of Kodaira–Ramanujam’s vanishing theorem*, Math. Ann. **261** (1982), 43–46.
- Kod53 K. Kodaira, *On a differential-geometric method in the theory of analytic stacks*, Proc. Natl. Acad. Sci. USA **39** (1953), 1268–1273.
- Kol86 J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. of Math. (2) **123** (1986), 11–42.
- Miy79 Y. Miyaoka, *On the Mumford–Ramanujam vanishing theorem on a surface*, in *Proceedings of Géométrie Algébrique d’Angers*, July 1979 (Sijthoff & Noordhoff, Alphen aan den Rijn Germantown, MD, 1980), 239–247.
- Mum67 D. Mumford, *Pathologies III*, Amer. J. Math. **89** (1967), 94–104.
- Ram72 C. P. Ramanujam, *Remarks on the Kodaira vanishing theorem*, J. Indian Math. Soc. (N.S.) (1972), 41–51.
- Ram74 C. P. Ramanujam, *Supplement to the article [Ram72]*, J. Indian Math. Soc. (N.S.) **38** (1974), 121–124.
- Ray78 M. Raynaud, *Contre-exemple au vanishing theorem en caractéristique  $p > 0$* , in *C. P. Ramanujam—A tribute*, Studies in Mathematics, vol. 8 (Tata Institute of Fundamental Research, Bombay, 1978), 273–278.
- Ser55 J.-P. Serre, *Faisceaux algébriques cohérents*, Ann. of Math. (2) **61** (1955), 197–278.
- Sho93 V. V. Shokurov, *Three-dimensional log perestroikas*, Russian Acad. Sci. Izv. Math. **40** (1993), 95–202.

# AN INJECTIVITY THEOREM

- Sho03 V. V. Shokurov, *Prelimiting flips*, in *Birational geometry: linear systems and finitely generated algebras: collected papers*, Proceedings of the Steklov Institute of Mathematics, vol. 240, eds V. A. Iskovskikh and V. V. Shokurov (Nauka, Moscow, 2003), 75–213.
- Tan71 S. G. Tankeev, *On  $n$ -dimensional canonically polarized varieties and varieties of fundamental type*, Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 31–44.
- Vie82 E. Viehweg, *Vanishing theorems*, J. Reine Angew. Math. **335** (1982), 1–8.
- Zar35 O. Zariski, *Algebraic surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 3 (Springer, 1935).

Florin Ambro [florin.ambro@imar.ro](mailto:florin.ambro@imar.ro)

Institute of Mathematics ‘Simion Stoilow’ of the Romanian Academy, PO box 1-764,  
RO-014700 Bucharest, Romania